
Hieracium macilentum agg.

Distribution





Map info

revised records

unrevised records

On the map are not visualized records without the coordinates and records marked as incorrect or doubtful.









Habitus and growth type

Height [m]: **0.2–0.4**
Growth form: **clonal herb**
Life form: **hemicryptophyte**

Leaf

Leaf presence and metamorphosis: **leaves present, not modified**
Leaf arrangement (phyllotaxis): **alternate, rosulate**
Leaf shape: **simple – entire**
Stipules: **absent**
Petiole: **both present and absent**

Flower

Flowering period [month]: **July–August**
Flower colour: **yellow**
Flower symmetry: **zygomorphic**
Perianth type: **calyx reduced, corolla present**
Perianth fusion: **fused**
Shape of the sympetalous corolla or syntepalous perianth: **ligulate**
Calyx fusion: **pappus**
Inflorescence type: **anthella ex anthodiis composita**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Generative reproduction type: **obligate apomixis**

Fruit, seed and dispersal

Fruit type: **dry fruit – achene/cypsela/samara**
Dispersal unit (diaspore): **fruit, infrutescence or its part**
Dispersal strategy: **Epilobium (mainly anemochory and autochory)**
Myrmecochory: **probably non-myrmecochorous**

Trophic mode

Parasitism and mycoheterotrophy: **autotrophic**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in the Czech Republic: **native**

Habitat and sociology

Occurrence in habitats

2 Alpine and subalpine grasslands

2B Subalpine tall-forb and tall-grass vegetation: **2 – optimum**

7 Acidophilous grasslands

7A Subalpine and montane acidophilous grasslands: **2 – optimum**

11 Heathlands and scrub

11A Dry lowland to subalpine heathlands: **2 – optimum**
11D Subalpine acidophilous Pinus mugo scrub: **2 – optimum**
11H Subalpine deciduous scrub: **1 – rare occurrence**

12 Forests

12R Acidophilous spruce forests: **1 – rare occurrence**

Affinity to the forest environment

Affinity to the forest environment in Thermophyticum: **0 – taxon that does not spontaneously occur in Czech forests**
Affinity to the forest environment in Mesophyticum and Oreophyticum: **0 – taxon that does not spontaneously occur in Czech forests**

Distribution and frequency

Elevational belt in the Czech Republic: **montane belt, subalpine belt**
Occurrence frequency in the basic grid mapping cells and quadrants of the basic grid mapping cells: 7
taxon.data.freq_in_quad: 7

Number of habitats with taxon occurrence in the Czech Republic

Number of narrow habitats in which the taxon occurs: **6**
Number of narrow habitats in which the taxon has its optimum: **4**
Number of broad habitats in which the taxon occurs: **4**
Number of broad habitats in which the taxon has its optimum: **3**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz