
Alchemilla filicaulis

Distribution





Map info

revised records

unrevised records

On the map are not visualized records without the coordinates and records marked as incorrect or doubtful.







Habitus and growth type

Height [m]: **0.1–0.25**
Growth form: **clonal herb**
Life form: **hemicryptophyte**
Life strategy: **CSR – competitor/stress-tolerator/ruderal**

Leaf

Leaf presence and metamorphosis: **leaves present, not modified**
Leaf arrangement (phyllotaxis): **rosulate**
Leaf shape: **simple – palmately divided**
Stipules: **present**
Petiole: **present**
Leaf life span: **summer green**

Flower

Flowering period [month]: **May–June**
Flower colour: **yellow-green**
Perianth type: **calyx present, corolla absent**
Calyx fusion: **hypanthium**
Inflorescence type: **racemus e cincinnis compositus**
Dicliny: **synoecious**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Generative reproduction type: **obligate apomixis**



Fruit, seed and dispersal

Fruit type: **dry fruit – achene/cypsela/samara**
Reproduction type: **by seed/spores and vegetatively**
Dispersal unit (diaspore): **fruit, infrutescence or its part**
Dispersal strategy: **Allium (mainly autochory)**
Myrmecochory: **non-myrmecochorous (b)**

Belowground organs and clonality



Shoot metamorphosis: **rhizome**
Storage organ: **rhizome**

Trophic mode

Parasitism and mycoheterotrophy: **autotrophic**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin



Origin in the Czech Republic: **native**

Habitat and sociology

Occurrence in habitats

4 Wetland and riverine herbaceous vegetation

4E Reed vegetation of brooks: **1 – rare occurrence**
4K Petasites fringes of montane brooks: **1 – rare occurrence**

5 Vegetation of springs and mires

5D Calcareous fens: **1 – rare occurrence**

6 Meadows and mesic pastures

6A Mesic Arrhenatherum meadows: **1 – rare occurrence**
6B Montane mesic meadows: **2 – optimum**
6C Pastures and park grasslands: **2 – optimum**
6D Alluvial meadows of lowland rivers: **1 – rare occurrence**
6E Wet Cirsium meadows: **2 – optimum**
6F Intermittently wet Molinia meadows: **2 – optimum**
6G Vegetation of wet disturbed soils: **1 – rare occurrence**



12 Forests

12U Plantations of broad-leaved non-native trees: **1 – rare occurrence**

Affinity to the forest environment

Affinity to the forest environment in Thermophyticum: **0 – taxon that does not spontaneously occur in Czech forests**
Affinity to the forest environment in Mesophyticum and Oreophyticum: **0 – taxon that does not spontaneously occur in Czech forests**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Distribution and frequency



Floristic zone: **arctic, boreal, northern temperate, submeridional, subtropical**
Floristic region: **Europe, Eastern America, Greenland**
Elevational belt in the Czech Republic: **colline belt, submontane belt**
Occurrence frequency in the basic grid mapping cells and quadrants of the basic grid mapping cells: 30
taxon.data.freq_in_quad: 36

Number of habitats with taxon occurrence in the Czech Republic

Number of narrow habitats in which the taxon occurs: **11**
Number of narrow habitats in which the taxon has its optimum: **4**
Number of broad habitats in which the taxon occurs: **4**
Number of broad habitats in which the taxon has its optimum: **1**



Threats and protection

Legal protection: **not protected by law**





© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz