
Alchemilla gruneica

Distribution



Habitus and growth type

Height [m]: **0.1–0.4**
Growth form: **clonal herb**
Life form: **hemicryptophyte**

Leaf

Leaf presence and metamorphosis: **leaves present, not modified**
Leaf arrangement (phyllotaxis): **rosulate**
Leaf shape: **simple – palmately divided**
Stipules: **present**
Petiole: **present**

Flower

Flowering period [month]: **May–July**
Flower colour: **yellow-green**
Perianth type: **calyx present, corolla absent**
Calyx fusion: **hypanthium**
Inflorescence type: **racemus e cincinnis compositus**
Generative reproduction type: **obligate apomixis**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Fruit, seed and dispersal

Fruit type: **dry fruit – achene/cypsela/samara**
Dispersal unit (diaspore): **fruit, infrutescence or its part**
Dispersal strategy: **Allium (mainly autochory)**
Myrmecochory: **non-myrmecochorous (b)**

Trophic mode

Parasitism and mycoheterotrophy: **autotrophic**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in the Czech Republic: **native**

Habitat and sociology

Occurrence in habitats

6 Meadows and mesic pastures

6A Mesic Arrhenatherum meadows: **1 – rare occurrence**
6B Montane mesic meadows: **2 – optimum**
6C Pastures and park grasslands: **2 – optimum**
6F Intermittently wet Molinia meadows: **1 – rare occurrence**
6G Vegetation of wet disturbed soils: **1 – rare occurrence**

7 Acidophilous grasslands

7A Subalpine and montane acidophilous grasslands: **2 – optimum**
7B Submontane Nardus grasslands: **1 – rare occurrence**

Affinity to the forest environment

Affinity to the forest environment in Thermophyticum: **0 – taxon that does not spontaneously occur in Czech forests**
Affinity to the forest environment in Mesophyticum and Oreophyticum: **0 – taxon that does not spontaneously occur in Czech forests**

Distribution and frequency

Elevational belt in the Czech Republic: **submontane belt, montane belt**
Occurrence frequency in the basic grid mapping cells and quadrants of the basic grid mapping cells: 24
taxon.data.freq_in_quad: 38

Number of habitats with taxon occurrence in the Czech Republic

Number of narrow habitats in which the taxon occurs: **7**
Number of narrow habitats in which the taxon has its optimum: **3**
Number of broad habitats in which the taxon occurs: **2**
Number of broad habitats in which the taxon has its optimum: **2**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Threats and protection

Red List 2017 (national categories): **C3 – vulnerable taxon**
Red List 2017 (IUCN categories): **LC – least concern**
Legal protection: **not protected by law**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz