
Amaranthus viridis

Distribution





Map info

revised records

unrevised records

On the map are not visualized records without the coordinates and records marked as incorrect or doubtful.







Habitus and growth type

Height [m]: **0.1–0.9**
Growth form: **annual herb**
Life form: **therophyte**

Leaf

Leaf presence and metamorphosis: **leaves present, not modified**
Leaf arrangement (phyllotaxis): **alternate**
Leaf shape: **simple – entire**
Stipules: **absent**
Petiole: **present**

Flower

Flowering period [month]: **August–October**
Flower colour: **green**
Flower symmetry: **actinomorphic**
Perianth type: **homochlamydeous**
Perianth fusion: **free**
Inflorescence type: **pseudospica**
Generative reproduction type: **mixed mating**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Fruit, seed and dispersal

Fruit type: **dry fruit – achene/cypsela/samara, dry fruit – capsule**
Dispersal unit (diaspore): **seed, fruit, infrutescence or its part**
Dispersal strategy: **Allium (mainly autochory)**
Myrmecochory: **non-myrmecochorous (b)**

Trophic mode

Parasitism and mycoheterotrophy: **autotrophic**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in the Czech Republic: **neophyte**
Invasion status: **casual**
Geographic origin: **South America**
Year of the first record in the wild: **1964**
Period of introduction: **Recent Past (1950–2000)**
Introduction pathway: **unintentional – industry**

Ecological indicator values

Ellenberg-type indicator values

Light indicator value: **8 – light plant, only exceptionally occurring at less than 40% of diffuse radiation incident in an open area**
Temperature indicator value: **8 – transition between values 7 and 9**
Moisture indicator value: **4 – transition between values 3 and 5**
Reaction indicator value: **7 – indicator of slightly acidic to slightly basic conditions, never occurring in very acidic conditions**
Nutrient indicator value: **8 – pronounced nutrient indicator**
Salinity indicator value: **1 – salt tolerant, mostly on low-salt to salt-free soils, but occasionally on slightly salty soils**

Habitat and sociology

Affinity to the forest environment

Affinity to the forest environment in Thermophyticum: **0 – taxon that does not spontaneously occur in Czech forests**
Affinity to the forest environment in Mesophyticum and Oreophyticum: **0 – taxon that does not spontaneously occur in Czech forests**

Distribution and frequency

Occurrence frequency in the basic grid mapping cells and quadrants of the basic grid mapping cells: 13
taxon.data.freq_in_quad: 15

Commonness in vegetation plots from the Czech Republic

Occurrence frequency in vegetation plots: **0 %**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Occurrence frequency in vegetation plots with a cover above 5%: **0 %**
Occurrence frequency in vegetation plots with a cover above 25%: **0 %**
Occurrence frequency in vegetation plots with a cover above 50%: **0 %**
Mean percentage cover in vegetation plots: **1.7 %**
Maximum percentage cover in vegetation plots: **2 %**

Threats and protection

Red List 2017 (national categories): **taxon is not on the Red List**
Red List 2017 (IUCN categories): **LC(NA) – least concern (taxon is not on the Red List)**
Legal protection: **not protected by law**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz