
Amelanchier ovalis



Distribution



Map info

- revised records
- unrevised records

On the map are not visualized records without the coordinates and records marked as incorrect or doubtful.







Habitus and growth type

Growth form: **shrub**
Life strategy: **C – competitor**
Life strategy (Pierce method based on leaf traits): **S/CSR**
Life strategy (Pierce method, C-score): **16.2 %**
Life strategy (Pierce method, S-score): **64.7 %**
Life strategy (Pierce method, R-score): **19 %**

Leaf

Leaf presence and metamorphosis: **leaves present, not modified**
Leaf arrangement (phyllotaxis): **alternate**
Leaf shape: **simple – entire**
Stipules: **present**
Petiole: **present**
Leaf life span: **summer green**
Leaf anatomy: **mesomorphic**
Functional leaf type in woody plants: **broad deciduous or semi-deciduous**

Flower

Flowering phase: **3 Prunus avium-Ranunculus auricomus (end of early spring)**
Flower colour: **white**


© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Flower symmetry: **actinomorphic**
Perianth type: **calyx and corolla**
Perianth fusion: **free**
Calyx fusion: **hypanthium**
Inflorescence type: **racemus**
Dicliny: **synoecious**
Pollination syndrome: **insect-pollination, selfing**



Fruit, seed and dispersal

Fruit type: **fleshy fruit – pome**
Fruit colour: **brown, black**
Reproduction type: **mostly by seed/spores, rarely vegetatively**
Myrmecochory: **non-myrmecochorous (b)**



Belowground organs and clonality

Root metamorphosis: **root shoot**
Position of root buds: **lateral roots**
Role of root buds in life-history of a plant: **regenerative**



Bud bank

Number of buds per shoot at the soil surface (root buds excluded): **0**
Number of buds per shoot at a depth of 0–10 cm (root buds excluded): **5**
Number of buds per shoot at a depth greater than 10 cm (root buds excluded): **5**
Size of the belowground bud bank (root buds excluded): **10**
Depth of the belowground bud bank (root buds exluded) [cm]: **10**
Number of buds per shoot at the soil surface (root buds included): **0**
Number of buds per shoot at a depth of 0–10 cm (root buds included): **20**
Number of buds per shoot at a depth greater than 10 cm (root buds included): **20**
Size of the belowground bud bank (root buds included): **40**
Depth of the belowground bud bank (root buds included) [cm]: **10**



Trophic mode

Parasitism and mycoheterotrophy: **autotrophic**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**



Taxon origin

Origin in the Czech Republic: **cultivated only**

Habitat and sociology

Affinity to the forest environment

Affinity to the forest environment in Thermophyticum: **0 – taxon that does not spontaneously occur in Czech forests**
Affinity to the forest environment in Mesophyticum and Oreophyticum: **0 – taxon that does not spontaneously occur in Czech forests**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Distribution and frequency



Floristic zone: **southern temperate, submeridional, meridional**
Floristic region: **Europe**
Occurrence frequency in the basic grid mapping cells and quadrants of the basic grid mapping cells: 22
taxon.data.freq_in_quad: 25

Threats and protection

Red List 2017 (national categories): **taxon is not on the Red List**
Red List 2017 (IUCN categories): **LC(NA) – least concern (taxon is not on the Red List)**
Legal protection: **not protected by law**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz