Atriplex patula

Distribution



Map info

revised records

unrevised records

On the map are not visualized records without the coordinates and records marked as incorrect or doubtful.

Habitus and growth type

Height [m]: **0.2–0.8**
Growth form: **annual herb**
Life form: **therophyte**
Life strategy: **CR – competitor/ruderal**
Life strategy (Pierce method based on leaf traits): **CR**
Life strategy (Pierce method, C-score): **53.9 %**
Life strategy (Pierce method, S-score): **0 %**
Life strategy (Pierce method, R-score): **46.1 %**

Leaf

Leaf presence and metamorphosis: **leaves present, not modified**
Leaf arrangement (phyllotaxis): **alternate, opposite**
Leaf shape: **simple – entire**
Stipules: **absent**
Petiole: **present**
Leaf life span: **summer green**
Leaf anatomy: **scleromorphic, mesomorphic**

Flower

Flowering period [month]: **July–October**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Flowering phase: **8 Clematis vitalba-Galium sylvaticum (mid-summer)**
Flower colour: **green**
Flower symmetry: **actinomorphic**
Perianth type: **homochlamydeous, sometimes absent**
Perianth fusion: **free**
Inflorescence type: **pseudospica**
Dicliny: **monoecious**
Generative reproduction type: **facultative allogamy**
Pollination syndrome: **wind-pollination, insect-pollination, selfing**

Fruit, seed and dispersal

Fruit type: **dry fruit – achene/cypsela/samara**
Fruit colour: **brown, black**
Reproduction type: **only by seed/spores**
Dispersal unit (diaspore): **fruit, infrutescence or its part**
Dispersal strategy: **Allium (mainly autochory)**
Myrmecochory: **non-myrmecochorous (b)**

Belowground organs and clonality

Shoot life span (cyclicity): **monocyclic shoots prevailing**
Primary root: **present**

Bud bank

Number of buds per shoot at the soil surface (root buds excluded): **3**
Number of buds per shoot at a depth of 0–10 cm (root buds excluded): **0**
Number of buds per shoot at a depth greater than 10 cm (root buds excluded): **0**
Size of the belowground bud bank (root buds excluded): **3**
Depth of the belowground bud bank (root buds exluded) [cm]: **1**
Number of buds per shoot at the soil surface (root buds included): **3**
Number of buds per shoot at a depth of 0–10 cm (root buds included): **0**
Number of buds per shoot at a depth greater than 10 cm (root buds included): **0**
Size of the belowground bud bank (root buds included): **3**
Depth of the belowground bud bank (root buds included) [cm]: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotrophic**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Karyology

Chromosome number (2n): **36**
Ploidy level (x): **4**
2C genome size [Mbp]: **3707.81**
1Cx monoploid genome size [Mbp]: **926.95**
Genomic GC content: **38.3 %**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Taxon origin

Origin in the Czech Republic: **archaeophyte**
Invasion status: **naturalized**
Geographic origin: **Europe, Mediterranean, Asia**
Period of introduction: **Neolithic (5600–4200 BCE)**
Introduction pathway: **unintentional – agriculture, unintentional – anthropogenic**

Ecological indicator values

Ellenberg-type indicator values

Light indicator value: **7 – half-light plant, mostly occurring at full light, but also in the shade up to about 30% of diffuse radiation incident in an open area**
Temperature indicator value: **6 – transition between values 5 and 7**
Moisture indicator value: **5 – indicator of fresh soils, focus on soils of average moisture, missing on wet and on soils that frequently dry out**
Reaction indicator value: **7 – indicator of slightly acidic to slightly basic conditions, never occurring in very acidic conditions**
Nutrient indicator value: **7 – occurring at nutrient-rich sites more often than at average sites and only exceptionally at poor sites**
Salinity indicator value: **1 – salt tolerant, mostly on low-salt to salt-free soils, but occasionally on slightly salty soils**

Indicator values for disturbance

Whole-community disturbance frequency indicator value: **0.02**
Herb layer disturbance frequency indicator value: **0.08**
Whole-community disturbance severity indicator value: **0.9**
Herb layer disturbance severity indicator value: **0.9**
Whole-community structure based disturbance indicator value: **0.67**
Herb layer structure-based disturbance indicator value: **0.74**

Habitat and sociology

Occurrence in habitats

4 Wetland and riverine herbaceous vegetation

4H Vegetation of low annual hygrophilous herbs: **1 – rare occurrence**
4I Vegetation of nitrophilous annual hygrophilous herbs: **2 – optimum**
4J River gravel banks: **1 – rare occurrence**
4L Nitrophilous herbaceous fringes of lowland rivers: **1 – rare occurrence**

6 Meadows and mesic pastures

6G Vegetation of wet disturbed soils: **1 – rare occurrence**

10 Saline vegetation

10I Inland saline meadows: **1 – rare occurrence**
10J Saline steppes: **1 – rare occurrence**

11 Heathlands and scrub

11L Tall mesic and xeric shrub: **1 – rare occurrence**
11R Scrub and pioneer woodland of forests clearings: **1 – rare occurrence**

13 Anthropogenic vegetation

13A Annual vegetation of ruderal habitats: **3 – dominant**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

13B Annual vegetation of arable land: **2 – optimum**
13C Annual vegetation of trampled habitats: **1 – rare occurrence**
13D Perennial thermophilous ruderal vegetation: **2 – optimum**
13E Perennial nitrophilous herbaceous vegetation of mesic sites: **1 – rare occurrence**

Affinity to the forest environment

Affinity to the forest environment in Thermophyticum: **0 – taxon that does not spontaneously occur in Czech forests**
Affinity to the forest environment in Mesophyticum and Oreophyticum: **0 – taxon that does not spontaneously occur in Czech forests**

Diagnostic taxon

Diagnostic taxon of classes: **XB *Stellarietea mediae***
Diagnostic taxon of alliances: **MBB *Chenopodion rubri***, **XBA *Caucalidion***, **XBG *Atriplicion***
Diagnostic taxon of associations: **MBB02 *Bidenti frondosae-Atriplicetum prostratae***, **XBA03 *Euphorbio exiguae-Melandrietum noctiflori***, **XBG02 *Chenopodietum urbici***, **XBG10 *Chamaeplietum officinalis***

Constant taxon

Constant taxon of associations: **MBB02 *Bidenti frondosae-Atriplicetum prostratae***, **XBA01 *Caucalido platycarpi-Conringietum orientalis***, **XBA03 *Euphorbio exiguae-Melandrietum noctiflori***, **XBG01 *Chenopodietum stricti***, **XBG02 *Chenopodietum urbici***, **XBG10 *Chamaeplietum officinalis***

Ecological specialization indices

Ecological specialization index for all vegetation types: **4.9**
Ecological specialization index for non-forest vegetation: **4.9**
Ecological specialization index for forest vegetation: **4.2**

Colonization ability

Index of colonization success (ICS): **5**
Index of colonization potential (ICP): **2**
Optimum successional age [years]: **4.5**

Distribution and frequency

Floristic zone: **boreal, northern temperate, southern temperate, submeridional, meridional**
Floristic region: **Europe, Western Asia, Siberia, Americas**
Distribution range extension along the continentality gradient: **6**
Elevational belt in the Czech Republic: **lowlands, colline belt, submontane belt**
Occurrence frequency in the basic grid mapping cells and quadrants of the basic grid mapping cells: 608
taxon.data.freq_in_quad: 1774

Commonness in vegetation plots from the Czech Republic

Occurrence frequency in vegetation plots: **2.8 %**
Occurrence frequency in vegetation plots with a cover above 5%: **7.9 %**
Occurrence frequency in vegetation plots with a cover above 25%: **1.8 %**
Occurrence frequency in vegetation plots with a cover above 50%: **0.2 %**
Mean percentage cover in vegetation plots: **3.6 %**
Maximum percentage cover in vegetation plots: **88 %**

Number of habitats with taxon occurrence in the Czech Republic

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Number of narrow habitats in which the taxon occurs: **14**
Number of narrow habitats in which the taxon has its optimum: **4**
Number of broad habitats in which the taxon occurs: **5**
Number of broad habitats in which the taxon has its optimum: **2**

Threats and protection

Red List 2017 (national categories): **taxon is not on the Red List**
Red List 2017 (IUCN categories): **LC(NA) – least concern (taxon is not on the Red List)**
Legal protection: **not protected by law**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz