


# *Dysphania schraderiana*

## Distribution



© Dana Michalčová

### Map info

● revised records

○ unrevised records

On the map are not visualized records without the coordinates and records marked as incorrect or doubtful.



© Dana Michalčová

## Habitus and growth type

Height [m]: **0.2-0.9**

Growth form: **annual herb**

Life form: **therophyte**

Life strategy: **CR - competitor/ruderal**

## Leaf

Leaf presence and metamorphosis: **leaves present, not modified**

Leaf arrangement (phyllotaxis): **alternate**

Leaf shape: **simple - entire**

Stipules: **absent**

Petiole: **present**

Leaf life span: **summer green**

Leaf anatomy: **mesomorphic**

## Flower

Flowering period [month]: **July-September**

Flower colour: **green**

Flower symmetry: **actinomorphic**


Perianth type: **homochlamydeous**

Perianth fusion: **fused**



© Dana Michalčová

Inflorescence type: **panicula e dichasiis composita**  
Dicliny: **synoecious**  
Generative reproduction type: **facultative allogamy**  
Pollination syndrome: **wind-pollination**



### **Fruit, seed and dispersal**

Fruit type: **dry fruit - achene/cypsela/samara**  
Fruit colour: **brown**  
Reproduction type: **only by seed/spores**  
Dispersal unit (diaspore): **seed, fruit, infrutescence or its part**  
Dispersal strategy: **Allium (mainly autochory)**  
Myrmecochory: **non-myrmecochorous (b)**

### **Trophic mode**

Parasitism and mycoheterotrophy: **autotrophic**  
Carnivory: **non-carnivorous**  
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

### **Taxon origin**

Origin in the Czech Republic: **neophyte**  
Invasion status: **casual**  
Geographic origin: **Africa**  
Year of the first record in the wild: **1894**  
Period of introduction: **Late Modern Period (1800-1950)**  
Introduction pathway: **intentional - crops**

### **Habitat and sociology**

Affinity to the forest environment  
Affinity to the forest environment in Thermophyticum: **0 - taxon that does not spontaneously occur in Czech forests**  
Affinity to the forest environment in Mesophyticum and Oreophyticum: **0 - taxon that does not spontaneously occur in Czech forests**

### **Distribution and frequency**

Floristic zone: **subtropical, austral or antarctic**  
Floristic region: **Africa**  
Elevational belt in the Czech Republic: **lowlands, colline belt**  
Occurrence frequency in the basic grid mapping cells and quadrants of the basic grid mapping cells: 16  
taxon.data.freq\_in\_quad: 17

### **Threats and protection**

Red List 2017 (national categories): **taxon is not on the Red List**  
Red List 2017 (IUCN categories): **LC(NA) - least concern (taxon is not on the Red List)**

**List)**

Legal protection: **not protected by law**