
Galatella linosyris

Distribution





Map info

revised records

unrevised records

On the map are not visualized records without the coordinates and records marked as incorrect or doubtful.





Habitus and growth type

Height [m]: **0.15–0.4**
Growth form: **clonal herb**
Life form: **hemicryptophyte**
Life strategy: **CSR – competitor/stress-tolerator/ruderal**
Life strategy (Pierce method based on leaf traits): **S/SR**
Life strategy (Pierce method, C-score): **6.9 %**
Life strategy (Pierce method, S-score): **78.2 %**
Life strategy (Pierce method, R-score): **15 %**

Leaf

Leaf presence and metamorphosis: **leaves present, not modified**
Leaf arrangement (phyllotaxis): **alternate**
Leaf shape: **simple – entire**
Stipules: **absent**
Petiole: **absent**
Leaf life span: **summer green**
Leaf anatomy: **scleromorphic, mesomorphic**

Flower

Flowering period [month]: **July–September**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Flowering phase: **8 Clematis vitalba-Galium sylvaticum (mid-summer)**
Flower colour: **yellow**
Flower symmetry: **actinomorphic**
Perianth type: **calyx reduced, corolla present**
Perianth fusion: **fused**
Shape of the sympetalous corolla or syntepalous perianth: **tubular**
Calyx fusion: **pappus**
Inflorescence type: **corymbothyrsus ex anthodiis compositus**
Dicliny: **synoecious**
Generative reproduction type: **allogamy self-incompatibility, facultative allogamy**
Pollination syndrome: **insect-pollination, geitonogamy**




Fruit, seed and dispersal

Fruit type: **dry fruit – achene/cypsela/samara**
Fruit colour: **brown**
Reproduction type: **by seed/spores and vegetatively**
Dispersal unit (diaspore): **fruit, infrutescence or its part**
Dispersal strategy: **Epilobium (mainly anemochory and autochory)**
Myrmecochory: **probably non-myrmecochorous nv**



Belowground organs and clonality

Shoot metamorphosis: **rhizome**
Storage organ: **rhizome**
Type of clonal growth organ: **hypogeogenous rhizome**
Freely dispersible organs of clonal growth: **absent**
Shoot life span (cyclicity): **monocyclic shoots prevailing**
Branching type of stem-derived organs of clonal growth: **sympodial**
Primary root: **absent**
Persistence of the clonal growth organ [year]: **2**
Number of clonal offspring: **2.7**
Lateral spreading distance by clonal growth [m]: **0.09**
Clonal index: **4**



Bud bank

Number of buds per shoot at the soil surface (root buds excluded): **5**
Number of buds per shoot at a depth of 0–10 cm (root buds excluded): **12**
Number of buds per shoot at a depth greater than 10 cm (root buds excluded): **0**
Size of the belowground bud bank (root buds excluded): **17**
Depth of the belowground bud bank (root buds exluded) [cm]: **4**
Number of buds per shoot at the soil surface (root buds included): **5**
Number of buds per shoot at a depth of 0–10 cm (root buds included): **12**
Number of buds per shoot at a depth greater than 10 cm (root buds included): **0**
Size of the belowground bud bank (root buds included): **17**
Depth of the belowground bud bank (root buds included) [cm]: **4**



Trophic mode

Parasitism and mycoheterotrophy: **autotrophic**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz



Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Karyology



Chromosome number (2n): **18 (36)**
Ploidy level (x): **2 (4)**
2C genome size [Mbp]: **8421.02**
1Cx monoploid genome size [Mbp]: **4210.51**
Genomic GC content: **39.3 %**

Taxon origin

Origin in the Czech Republic: **native**



Ecological indicator values

Ellenberg-type indicator values

Light indicator value: **8 – light plant, only exceptionally occurring at less than 40% of diffuse radiation incident in an open area**
Temperature indicator value: **7 – heat indicator, occurring in relatively warm lowlands**
Moisture indicator value: **2 – transition between values 1 and 3**
Reaction indicator value: **8 – transition between values 7 and 9, occurring mostly in calcium-rich conditions**
Nutrient indicator value: **3 – occurring at nutrient-poor sites more frequently than at average sites and exceptionally at rich sites**
Salinity indicator value: **0 – not salt tolerant, glycophyte**



Indicator values for disturbance

Whole-community disturbance frequency indicator value: **-0.28**
Herb layer disturbance frequency indicator value: **-0.28**
Whole-community disturbance severity indicator value: **0.2**
Herb layer disturbance severity indicator value: **0.23**
Whole-community structure based disturbance indicator value: **0.49**
Herb layer structure-based disturbance indicator value: **0.6**



Habitat and sociology

Occurrence in habitats



1 Vegetation of cliffs, screes and walls


1D Mobile calcareous screes: **1 – rare occurrence**

8 Dry grasslands

8A Hercynian dry grasslands on rock outcrops: **2 – optimum**
8B Submediterranean dry grasslands on rock outcrops: **1 – rare occurrence**
8C Narrow-leaved sub-continental steppes: **2 – optimum**
8D Broad-leaved dry grasslands: **2 – optimum**
8F Thermophilous forest fringe vegetation: **2 – optimum**

11 Heathlands and scrub











11L Tall mesic and xeric shrub: **1 – rare occurrence**
11N Low xeric scrub: **2 – optimum**

12 Forests

12H Peri-Alpidic basiphilous thermophilous oak forests: **1 – rare occurrence**
12W Pine and larch plantations: **1 – rare occurrence**

Affinity to the forest environment

Affinity to the forest environment in Thermophyticum: **2.2 – taxon occurring partly in the forest, but mainly in open vegetation**

Diagnostic taxon

Diagnostic taxon of classes: **TH *Festuco-Brometea***
Diagnostic taxon of alliances: **THE *Cirsio-Brachypodion pinnati***
Diagnostic taxon of associations: **THD05 *Stipetum tirsae***, **THD06 *Astragalo exscapi-Crambetum tatariae***, **THE03 *Polygalo majoris-Brachypodietum pinnati***

Constant taxon

Constant taxon of associations: **THE03 *Polygalo majoris-Brachypodietum pinnati***

Ecological specialization indices

Ecological specialization index for all vegetation types: **5.4**
Ecological specialization index for non-forest vegetation: **5.4**
Ecological specialization index for forest vegetation: **5.4**

Colonization ability

Index of colonization success (ICS): **3**
Index of colonization potential (ICP): **1**
Optimum successional age [years]: **75**

Distribution and frequency

Floristic zone: **northern temperate, southern temperate, submeridional**
Floristic region: **Europe**
Continentality degree: **6**
Distribution range extension along the continentality gradient: **5**
Elevational belt in the Czech Republic: **lowlands, colline belt**
Occurrence frequency in the basic grid mapping cells and quadrants of the basic grid mapping cells: 100
taxon.data.freq_in_quad: 217

Commonness in vegetation plots from the Czech Republic

Occurrence frequency in vegetation plots: **0.6 %**
Occurrence frequency in vegetation plots with a cover above 5%: **17.6 %**
Occurrence frequency in vegetation plots with a cover above 25%: **3.6 %**
Occurrence frequency in vegetation plots with a cover above 50%: **0 %**
Mean percentage cover in vegetation plots: **5.2 %**
Maximum percentage cover in vegetation plots: **38 %**

Number of habitats with taxon occurrence in the Czech Republic


Number of narrow habitats in which the taxon occurs: **10**
Number of narrow habitats in which the taxon has its optimum: **5**
Number of broad habitats in which the taxon occurs: **4**
Number of broad habitats in which the taxon has its optimum: **2**

© 2014-2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Threats and protection

Red List 2017 (national categories): **C3 – vulnerable taxon**
Red List 2017 (IUCN categories): **NT – near threatened**
Legal protection: **vulnerable taxon**





© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz