
Gladiolus communis

Distribution



Habitus and growth type

Growth form: **clonal herb**

Leaf

Leaf presence and metamorphosis: **leaves present, not modified**
Leaf arrangement (phyllotaxis): **alternate**
Leaf shape: **simple – entire**
Stipules: **absent**
Petiole: **absent**

Flower

Flower colour: **pink-violet**
Flower symmetry: **zygomorphic**
Perianth type: **homochlamydeous**
Perianth fusion: **fused**
Shape of the sympetalous corolla or syntepalous perianth: **special type**
Inflorescence type: **racemus**

Fruit, seed and dispersal

Fruit type: **dry fruit – capsule**
Myrmecochory: **myrmecochorous nv**

© 2014-2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Belowground organs and clonality

Type of clonal growth organ: **belowground stem tuber**
Freely dispersible organs of clonal growth: **absent**
Shoot life span (cyclicity): **monocyclic shoots prevailing**
Branching type of stem-derived organs of clonal growth: **sympodial**
Primary root: **absent**
Persistence of the clonal growth organ [year]: **1**
Number of clonal offspring: **3.5**
Lateral spreading distance by clonal growth [m]: **0.01**
Clonal index: **3**

Bud bank

Number of buds per shoot at the soil surface (root buds excluded): **2**
Number of buds per shoot at a depth of 0–10 cm (root buds excluded): **5**
Number of buds per shoot at a depth greater than 10 cm (root buds excluded): **0**
Size of the belowground bud bank (root buds excluded): **7**
Depth of the belowground bud bank (root buds exluded) [cm]: **4**
Number of buds per shoot at the soil surface (root buds included): **2**
Number of buds per shoot at a depth of 0–10 cm (root buds included): **5**
Number of buds per shoot at a depth greater than 10 cm (root buds included): **0**
Size of the belowground bud bank (root buds included): **7**
Depth of the belowground bud bank (root buds included) [cm]: **4**

Trophic mode

Parasitism and mycoheterotrophy: **autotrophic**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in the Czech Republic: **cultivated only**

Habitat and sociology

Affinity to the forest environment

Affinity to the forest environment in Thermophyticum: **0 – taxon that does not spontaneously occur in Czech forests**
Affinity to the forest environment in Mesophyticum and Oreophyticum: **0 – taxon that does not spontaneously occur in Czech forests**

Threats and protection

Red List 2017 (national categories): **taxon is not on the Red List**
Red List 2017 (IUCN categories): **LC(NA) – least concern (taxon is not on the Red List)**
Legal protection: **not protected by law**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz