Herminium monorchis

Distribution



Map info

revised records

unrevised records

On the map are not visualized records without the coordinates and records marked as incorrect or doubtful.

Habitus and growth type

Height [m]: **0.08–0.25**
Growth form: **clonal herb**
Life form: **geophyte**
Life strategy: **CSR – competitor/stress-tolerator/ruderal**

Leaf

Leaf presence and metamorphosis: **leaves present, not modified**
Leaf arrangement (phyllotaxis): **alternate**
Leaf shape: **simple – entire**
Stipules: **absent**
Petiole: **absent**
Leaf life span: **summer green**
Leaf anatomy: **mesomorphic**

Flower

Flowering period [month]: **May–July**
Flower colour: **white**
Flower symmetry: **zygomorphic**
Perianth type: **homochlamydeous**
Perianth fusion: **free**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Inflorescence type: **spica**
Dicliny: **synoecious**
Generative reproduction type: **mixed mating**
Pollination syndrome: **insect-pollination, selfing**

Fruit, seed and dispersal

Fruit type: **dry fruit – capsule**
Fruit colour: **green, brown**
Reproduction type: **by seed/spores and vegetatively**
Dispersal unit (diaspore): **seed**
Dispersal strategy: **Lycopodium (mainly anemochory)**
Myrmecochory: **non-myrmecochorous (b)**

Belowground organs and clonality

Shoot metamorphosis: **stolon, stolon with tuberous tip**
Root metamorphosis: **root tuber**
Storage organ: **stolon, stolon with tuberous tip, root tuber**
Type of clonal growth organ: **hypogeogenous rhizome**
Freely dispersible organs of clonal growth: **absent**
Shoot life span (cyclicity): **monocyclic shoots prevailing**
Branching type of stem-derived organs of clonal growth: **sympodial**
Primary root: **absent**
Persistence of the clonal growth organ [year]: **1**
Number of clonal offspring: **1**
Lateral spreading distance by clonal growth [m]: **0.09**
Clonal index: **4**

Bud bank

Number of buds per shoot at the soil surface (root buds excluded): **5**
Number of buds per shoot at a depth of 0–10 cm (root buds excluded): **5**
Number of buds per shoot at a depth greater than 10 cm (root buds excluded): **0**
Size of the belowground bud bank (root buds excluded): **10**
Depth of the belowground bud bank (root buds exluded) [cm]: **3**
Number of buds per shoot at the soil surface (root buds included): **5**
Number of buds per shoot at a depth of 0–10 cm (root buds included): **5**
Number of buds per shoot at a depth greater than 10 cm (root buds included): **0**
Size of the belowground bud bank (root buds included): **10**
Depth of the belowground bud bank (root buds included) [cm]: **3**

Trophic mode

Parasitism and mycoheterotrophy: **partial or initial mycoheterotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in the Czech Republic: **native**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Ecological indicator values

Ellenberg-type indicator values

Light indicator value: **7 – half-light plant, mostly occurring at full light, but also in the shade up to about 30% of diffuse radiation incident in an open area**
Temperature indicator value: **5 – moderate heat indicator, occurring from lowland to montane belt, mainly in submontane-temperate areas**
Moisture indicator value: **5 – indicator of fresh soils, focus on soils of average moisture, missing on wet and on soils that frequently dry out**
Reaction indicator value: **8 – transition between values 7 and 9, occurring mostly in calcium-rich conditions**
Nutrient indicator value: **2 – transition between values 1 and 3**
Salinity indicator value: **0 – not salt tolerant, glycophyte**

Habitat and sociology

Affinity to the forest environment

Affinity to the forest environment in Thermophyticum: **0 – taxon that does not spontaneously occur in Czech forests**
Affinity to the forest environment in Mesophyticum and Oreophyticum: **0 – taxon that does not spontaneously occur in Czech forests**

Distribution and frequency

Floristic zone: **northern temperate, southern temperate, submeridional, meridional**
Floristic region: **Europe, Asia**
Continentality degree: **6**
Distribution range extension along the continentality gradient: **5**
Elevational belt in the Czech Republic: **colline belt, submontane belt**
Occurrence frequency in the basic grid mapping cells and quadrants of the basic grid mapping cells: 9
taxon.data.freq_in_quad: 12

Threats and protection

Red List 2017 (national categories): **A1 – extinct taxon**
Red List 2017 (IUCN categories): **RE – regionally extinct**
Legal protection: **critically threatened taxon**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz