
Hordeum secalinum

Distribution






Map info

revised records

unrevised records

On the map are not visualized records without the coordinates and records marked as incorrect or doubtful.





Habitus and growth type

Height [m]: **0.2–0.7**
Growth form: **clonal herb**
Life form: **hemicryptophyte**
Life strategy: **S – stress-tolerator**
Life strategy (Pierce method based on leaf traits): **R/CSR**
Life strategy (Pierce method, C-score): **16 %**
Life strategy (Pierce method, S-score): **32.8 %**
Life strategy (Pierce method, R-score): **51.2 %**

Leaf

Leaf presence and metamorphosis: **leaves present, not modified**
Leaf arrangement (phyllotaxis): **alternate**
Leaf shape: **simple – entire**
Stipules: **absent**
Petiole: **absent**
Leaf life span: **summer green**
Leaf anatomy: **mesomorphic**

Flower

Flowering period [month]: **June–July**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz



Flowering phase: **7 Ligustrum vulgare-Stachys sylvatica (end of early summer)**
Flower colour: **green**
Perianth type: **reduced**
Perianth fusion: **reduced**
Inflorescence type: **spica e spiculis composita**
Dicliny: **andromonoecious**
Pollination syndrome: **wind-pollination**

Fruit, seed and dispersal

Fruit type: **dry fruit – caryopsis**
Reproduction type: **only by seed/spores**
Myrmecochory: **non-myrmecochorous (b)**

Belowground organs and clonality

Storage organ: **tuft**
Type of clonal growth organ: **epigeogenous rhizome**
Freely dispersible organs of clonal growth: **absent**
Shoot life span (cyclicity): **dicyclic or polycyclic shoots prevailing**
Branching type of stem-derived organs of clonal growth: **sympodial**
Primary root: **absent**
Persistence of the clonal growth organ [year]: **4**
Number of clonal offspring: **3.5**
Lateral spreading distance by clonal growth [m]: **0.01**
Clonal index: **4**

Bud bank

Number of buds per shoot at the soil surface (root buds excluded): **5**
Number of buds per shoot at a depth of 0–10 cm (root buds excluded): **10**
Number of buds per shoot at a depth greater than 10 cm (root buds excluded): **0**
Size of the belowground bud bank (root buds excluded): **15**
Depth of the belowground bud bank (root buds exluded) [cm]: **4**
Number of buds per shoot at the soil surface (root buds included): **5**
Number of buds per shoot at a depth of 0–10 cm (root buds included): **10**
Number of buds per shoot at a depth greater than 10 cm (root buds included): **0**
Size of the belowground bud bank (root buds included): **15**
Depth of the belowground bud bank (root buds included) [cm]: **4**

Trophic mode

Parasitism and mycoheterotrophy: **autotrophic**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in the Czech Republic: **neophyte**
Invasion status: **casual**
Geographic origin: **Mediterranean**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Year of the first record in the wild: **1929**
Year of the last record: **1964**
Period of introduction: **Late Modern Period (1800–1950)**
Introduction pathway: **unintentional – agriculture, unintentional – industry, unintentional – anthropogenic**

Habitat and sociology

Affinity to the forest environment

Affinity to the forest environment in Thermophyticum: **0 – taxon that does not spontaneously occur in Czech forests**
Affinity to the forest environment in Mesophyticum and Oreophyticum: **0 – taxon that does not spontaneously occur in Czech forests**

Distribution and frequency

Floristic zone: **northern temperate, southern temperate, submeridional**
Floristic region: **Europe**
Occurrence frequency in the basic grid mapping cells and quadrants of the basic grid mapping cells: 5
taxon.data.freq_in_quad: 5

Commonness in vegetation plots from the Czech Republic

Occurrence frequency in vegetation plots: **0 %**
Occurrence frequency in vegetation plots with a cover above 5%: **0 %**
Occurrence frequency in vegetation plots with a cover above 25%: **0 %**
Occurrence frequency in vegetation plots with a cover above 50%: **0 %**
Mean percentage cover in vegetation plots: **3 %**
Maximum percentage cover in vegetation plots: **3 %**

Threats and protection

Red List 2017 (national categories): **taxon is not on the Red List**
Red List 2017 (IUCN categories): **LC(NA) – least concern (taxon is not on the Red List)**
Legal protection: **not protected by law**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz