
Chenopodium hybridum

Distribution





Map info

revised records

unrevised records

On the map are not visualized records without the coordinates and records marked as incorrect or doubtful.





Habitus and growth type

Height [m]: **0.4–1**
Growth form: **annual herb**
Life form: **therophyte**
Life strategy: **CR – competitor/ruderal**
Life strategy (Pierce method based on leaf traits): **CR**
Life strategy (Pierce method, C-score): **43.1 %**
Life strategy (Pierce method, S-score): **0 %**
Life strategy (Pierce method, R-score): **56.9 %**

Leaf

Leaf presence and metamorphosis: **leaves present, not modified**
Leaf arrangement (phyllotaxis): **alternate, opposite**
Leaf shape: **simple – entire**
Stipules: **absent**
Petiole: **present**
Leaf life span: **summer green**
Leaf anatomy: **mesomorphic**

Flower

Flowering period [month]: **June–October**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Flowering phase: **8 Clematis vitalba-Galium sylvaticum (mid-summer)**
Flower colour: **green**
Flower symmetry: **actinomorphic**
Perianth type: **homochlamydeous**
Perianth fusion: **free, fused**
Inflorescence type: **panicula e dichasiis composita**
Dicliny: **synoecious**
Generative reproduction type: **facultative allogamy**
Pollination syndrome: **wind-pollination**



Fruit, seed and dispersal

Fruit type: **dry fruit – achene/cypsela/samara**
Fruit colour: **brown, grey, black**
Reproduction type: **only by seed/spores**
Dispersal unit (diaspore): **seed, fruit, infrutescence or its part**
Dispersal strategy: **Allium (mainly autochory)**
Myrmecochory: **non-myrmecochorous (b)**



Belowground organs and clonality

Shoot life span (cyclicity): **monocyclic shoots prevailing**
Primary root: **present**

Bud bank

Number of buds per shoot at the soil surface (root buds excluded): **5**
Number of buds per shoot at a depth of 0–10 cm (root buds excluded): **0**
Number of buds per shoot at a depth greater than 10 cm (root buds excluded): **0**
Size of the belowground bud bank (root buds excluded): **5**
Depth of the belowground bud bank (root buds exluded) [cm]: **1**
Number of buds per shoot at the soil surface (root buds included): **5**
Number of buds per shoot at a depth of 0–10 cm (root buds included): **0**
Number of buds per shoot at a depth greater than 10 cm (root buds included): **0**
Size of the belowground bud bank (root buds included): **5**
Depth of the belowground bud bank (root buds included) [cm]: **1**



Trophic mode

Parasitism and mycoheterotrophy: **autotrophic**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Karyology

Chromosome number (2n): **18**
Ploidy level (x): **2**
2C genome size [Mbp]: **1045.59**
1Cx monoploid genome size [Mbp]: **522.8**
Genomic GC content: **39.6 %**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Taxon origin

Origin in the Czech Republic: **native**

Ecological indicator values

Ellenberg-type indicator values

Light indicator value: **7 – half-light plant, mostly occurring at full light, but also in the shade up to about 30% of diffuse radiation incident in an open area**
Temperature indicator value: **6 – transition between values 5 and 7**
Moisture indicator value: **5 – indicator of fresh soils, focus on soils of average moisture, missing on wet and on soils that frequently dry out**
Reaction indicator value: **7 – indicator of slightly acidic to slightly basic conditions, never occurring in very acidic conditions**
Nutrient indicator value: **8 – pronounced nutrient indicator**
Salinity indicator value: **2 – oligohaline, often on soils with very low salt content**

Indicator values for disturbance

Whole-community disturbance frequency indicator value: **-0.25**
Herb layer disturbance frequency indicator value: **-0.13**
Whole-community disturbance severity indicator value: **0.93**
Herb layer disturbance severity indicator value: **0.93**
Whole-community structure based disturbance indicator value: **0.65**
Herb layer structure-based disturbance indicator value: **0.66**

Habitat and sociology

Occurrence in habitats

1 Vegetation of cliffs, screes and walls

1A Calcareous cliffs: **1 – rare occurrence**
1D Mobile calcareous screes: **1 – rare occurrence**

4 Wetland and riverine herbaceous vegetation

4B Halophilous reed and sedge beds: **1 – rare occurrence**
4D Riverine reed vegetation: **1 – rare occurrence**
4I Vegetation of nitrophilous annual hygrophilous herbs: **1 – rare occurrence**
4J River gravel banks: **1 – rare occurrence**
4K Petasites fringes of montane brooks: **1 – rare occurrence**
4L Nitrophilous herbaceous fringes of lowland rivers: **1 – rare occurrence**

10 Saline vegetation

10I Inland saline meadows: **1 – rare occurrence**

11 Heathlands and scrub

11J Willow galleries of loamy and sandy river banks: **1 – rare occurrence**
11R Scrub and pioneer woodland of forests clearings: **1 – rare occurrence**

12 Forests

12T Robinia pseudacacia plantations: **1 – rare occurrence**
12U Plantations of broad-leaved non-native trees: **1 – rare occurrence**
12V Spruce plantations: **1 – rare occurrence**
12W Pine and larch plantations: **1 – rare occurrence**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

13 Anthropogenic vegetation

13A Annual vegetation of ruderal habitats: **2 – optimum**
13B Annual vegetation of arable land: **2 – optimum**
13C Annual vegetation of trampled habitats: **1 – rare occurrence**
13D Perennial thermophilous ruderal vegetation: **1 – rare occurrence**
13E Perennial nitrophilous herbaceous vegetation of mesic sites: **1 – rare occurrence**
13F Herbaceous vegetation of forests clearings and Rubus scrub: **1 – rare occurrence**

Affinity to the forest environment

Affinity to the forest environment in Thermophyticum: **0 – taxon that does not spontaneously occur in Czech forests**
Affinity to the forest environment in Mesophyticum and Oreophyticum: **0 – taxon that does not spontaneously occur in Czech forests**

Diagnostic taxon

Diagnostic taxon of alliances: **XBB *Veronico-Euphorbion***
Diagnostic taxon of associations: **XBB01 *Mercurialietum annuae***

Constant taxon

Constant taxon of associations: **XBB01 *Mercurialietum annuae***

Ecological specialization indices

Ecological specialization index for all vegetation types: **4.9**
Ecological specialization index for non-forest vegetation: **5**
Ecological specialization index for forest vegetation: **4.8**

Colonization ability

Index of colonization success (ICS): **1**
Index of colonization potential (ICP): **1**

Distribution and frequency

Floristic zone: **northern temperate, southern temperate, submeridional, meridional**
Floristic region: **Europe, Asia**
Continentality degree: **7**
Distribution range extension along the continentality gradient: **8**
Elevational belt in the Czech Republic: **lowlands, colline belt, submontane belt**
Occurrence frequency in the basic grid mapping cells and quadrants of the basic grid mapping cells: 474
taxon.data.freq_in_quad: 1152

Commonness in vegetation plots from the Czech Republic

Occurrence frequency in vegetation plots: **0.8 %**
Occurrence frequency in vegetation plots with a cover above 5%: **6.9 %**
Occurrence frequency in vegetation plots with a cover above 25%: **1.7 %**
Occurrence frequency in vegetation plots with a cover above 50%: **0.4 %**
Mean percentage cover in vegetation plots: **3.3 %**
Maximum percentage cover in vegetation plots: **88 %**

Number of habitats with taxon occurrence in the Czech Republic

Number of narrow habitats in which the taxon occurs: **21**
Number of narrow habitats in which the taxon has its optimum: **2**
Number of broad habitats in which the taxon occurs: **6**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Number of broad habitats in which the taxon has its optimum: **1**

Threats and protection

Red List 2017 (national categories): **taxon is not on the Red List**
Red List 2017 (IUCN categories): **LC(NA) – least concern (taxon is not on the Red List)**
Legal protection: **not protected by law**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz