
Iris pumila

Distribution





Map info

revised records

unrevised records

On the map are not visualized records without the coordinates and records marked as incorrect or doubtful.







Habitus and growth type

Height [m]: **0.1–0.15**
Growth form: **clonal herb**
Life form: **hemicryptophyte**
Life strategy: **CSR – competitor/stress-tolerator/ruderal**

Leaf

Leaf presence and metamorphosis: **leaves present, not modified**
Leaf arrangement (phyllotaxis): **alternate**
Leaf shape: **simple – entire**
Stipules: **absent**
Petiole: **absent**
Leaf life span: **summer green**
Leaf anatomy: **mesomorphic**

Flower

Flowering period [month]: **April–May**
Flower colour: **yellow-white, blue-violet**
Flower symmetry: **actinomorphic**
Perianth type: **homochlamydeous**
Perianth fusion: **fused**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Shape of the sympetalous corolla or syntepalous perianth: **special type**
Inflorescence type: **rhipidium**
Dicliny: **synoecious**
Generative reproduction type: **allogamy self-incompatibility**
Pollination syndrome: **insect-pollination**



Fruit, seed and dispersal

Fruit type: **dry fruit – capsule**
Fruit colour: **brown**
Reproduction type: **by seed/spores and vegetatively**
Dispersal unit (diaspore): **seed**
Dispersal strategy: **Allium (mainly autochory)**
Myrmecochory: **non-myrmecochorous (a) nv**



Belowground organs and clonality

Shoot metamorphosis: **rhizome**
Storage organ: **rhizome**
Type of clonal growth organ: **epigeogenous rhizome**
Freely dispersible organs of clonal growth: **absent**
Shoot life span (cyclicity): **dicyclic or polycyclic shoots prevailing**
Branching type of stem-derived organs of clonal growth: **sympodial**
Primary root: **absent**
Persistence of the clonal growth organ [year]: **4**
Number of clonal offspring: **0.9**
Lateral spreading distance by clonal growth [m]: **0.07**
Clonal index: **4**



Bud bank

Number of buds per shoot at the soil surface (root buds excluded): **5**
Number of buds per shoot at a depth of 0–10 cm (root buds excluded): **15**
Number of buds per shoot at a depth greater than 10 cm (root buds excluded): **0**
Size of the belowground bud bank (root buds excluded): **20**
Depth of the belowground bud bank (root buds exluded) [cm]: **4**
Number of buds per shoot at the soil surface (root buds included): **5**
Number of buds per shoot at a depth of 0–10 cm (root buds included): **15**
Number of buds per shoot at a depth greater than 10 cm (root buds included): **0**
Size of the belowground bud bank (root buds included): **20**
Depth of the belowground bud bank (root buds included) [cm]: **4**







Trophic mode

Parasitism and mycoheterotrophy: **autotrophic**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Karyology

Chromosome number (2n): **32**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz


Ploidy level (x): **4**
2C genome size [Mbp]: **20844.5**
1Cx monoploid genome size [Mbp]: **5211.13**
Genomic GC content: **41.7 %**



Taxon origin

Origin in the Czech Republic: **native**

Ecological indicator values



Ellenberg-type indicator values

Light indicator value: **9 – full light plant, occurring only in fully irradiated places, not at less than 50% of diffuse radiation incident in an open area**
Temperature indicator value: **8 – transition between values 7 and 9**
Moisture indicator value: **2 – transition between values 1 and 3**
Reaction indicator value: **9 – base and lime indicator, always occurring in calcium-rich conditions**
Nutrient indicator value: **2 – transition between values 1 and 3**
Salinity indicator value: **0 – not salt tolerant, glycophyte**

Indicator values for disturbance

Whole-community disturbance frequency indicator value: **-1.07**
Herb layer disturbance frequency indicator value: **-0.46**
Whole-community disturbance severity indicator value: **0.25**
Herb layer disturbance severity indicator value: **0.27**
Whole-community structure based disturbance indicator value: **0.62**
Herb layer structure-based disturbance indicator value: **0.64**



Habitat and sociology

Occurrence in habitats

8 Dry grasslands

8A Hercynian dry grasslands on rock outcrops: **1 – rare occurrence**
8B Submediterranean dry grasslands on rock outcrops: **2 – optimum**
8C Narrow-leaved sub-continental steppes: **2 – optimum**

9 Sand grasslands and rock-outcrop vegetation

9F Basiphilous vegetation of spring therophytes and succulents: **2 – optimum**

12 Forests

12H Peri-Alpidic basiphilous thermophilous oak forests: **2 – optimum**

Affinity to the forest environment

Affinity to the forest environment in Thermophyticum: **0 – taxon that does not spontaneously occur in Czech forests**
Affinity to the forest environment in Mesophyticum and Oreophyticum: **0 – taxon that does not spontaneously occur in Czech forests**

Diagnostic taxon

Diagnostic taxon of alliances: **TFF *Alysso alyssoidis-Sedion*, THB *Bromo pannonici-Festucion pallentis***
Diagnostic taxon of associations: **LCA02 *Lithospermo purpurocaerulei-Quercetum***

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

pubescentis*, TFF02 *Alysso alyssoidis-Sedetum*, THB01 *Poo badensis-Festucetum pallentis*, THD06 *Astragalo exscapi-Crambetum tatariae



Constant taxon

Constant taxon of associations: **THD06 *Astragalo exscapi-Crambetum tatariae***

Ecological specialization indices

Ecological specialization index for all vegetation types: **5.6**
Ecological specialization index for non-forest vegetation: **5.6**

Colonization ability

Index of colonization success (ICS): **2**
Index of colonization potential (ICP): **3**
Optimum successional age [years]: **18**

Distribution and frequency



Floristic zone: **submeridional**
Floristic region: **Europe**
Elevational belt in the Czech Republic: **colline belt**
Occurrence frequency in the basic grid mapping cells and quadrants of the basic grid mapping cells: 34
taxon.data.freq_in_quad: 58

Commonness in vegetation plots from the Czech Republic

Occurrence frequency in vegetation plots: **0.1 %**
Occurrence frequency in vegetation plots with a cover above 5%: **3.1 %**
Occurrence frequency in vegetation plots with a cover above 25%: **0 %**
Occurrence frequency in vegetation plots with a cover above 50%: **0 %**
Mean percentage cover in vegetation plots: **2.5 %**
Maximum percentage cover in vegetation plots: **13 %**

Number of habitats with taxon occurrence in the Czech Republic

Number of narrow habitats in which the taxon occurs: **5**
Number of narrow habitats in which the taxon has its optimum: **4**
Number of broad habitats in which the taxon occurs: **3**
Number of broad habitats in which the taxon has its optimum: **3**



Threats and protection

Red List 2017 (national categories): **C2r – endangered taxon, rare**
Red List 2017 (IUCN categories): **VU – vulnerable**
Legal protection: **endangered taxon**

