


Oreojuncus trifidus

Distribution



© Dana Michalčová

Map info

● revised records

○ unrevised records

On the map are not visualized records without the coordinates and records marked as incorrect or doubtful.



© Karel Fajmon



© 01112025



© Gianmaria Bonari

Habitus and growth type

Height [m]: **0.1-0.3**

Growth form: **clonal herb**

Life form: **hemicryptophyte**

Life strategy: **CSR - competitor/stress-tolerator/ruderal**

Life strategy (Pierce method based on leaf traits): **S/SR**

Life strategy (Pierce method, C-score): **6.2 %**

Life strategy (Pierce method, S-score): **75.4 %**

Life strategy (Pierce method, R-score): **18.5 %**

Leaf

Leaf presence and metamorphosis: **leaves present, not modified**

Leaf arrangement (phyllotaxis): **alternate**

Leaf shape: **simple - entire**

Stipules: **absent**

Petiole: **absent**

Leaf life span: **evergreen**

Leaf anatomy: **scleromorphic**

Flower

Flowering period [month]: **July-September**

Flower colour: **green**
Flower symmetry: **actinomorphic**
Perianth type: **homochlamydeous**
Perianth fusion: **free**
Inflorescence type: **anthella**
Dicliny: **synoecious**
Generative reproduction type: **facultative allogamy**
Pollination syndrome: **wind-pollination**



Fruit, seed and dispersal

Fruit type: **dry fruit - capsule**
Reproduction type: **by seed/spores and vegetatively**
Dispersal unit (diaspore): **seed**
Dispersal strategy: **Allium (mainly autochory)**
Myrmecochory: **probably non-myrmecochorous**

Belowground organs and clonality

Shoot metamorphosis: **rhizome**
Storage organ: **rhizome, tuft**
Type of clonal growth organ: **hypogeogenous rhizome**
Freely dispersible organs of clonal growth: **absent**
Shoot life span (cyclicality): **monocyclic shoots prevailing**
Branching type of stem-derived organs of clonal growth: **sympodial**
Primary root: **absent**
Persistence of the clonal growth organ [year]: **4**
Number of clonal offspring: **3.5**
Lateral spreading distance by clonal growth [m]: **0.08**
Clonal index: **5**

Bud bank

Number of buds per shoot at the soil surface (root buds excluded): **4**
Number of buds per shoot at a depth of 0–10 cm (root buds excluded): **15**
Number of buds per shoot at a depth greater than 10 cm (root buds excluded): **0**
Size of the belowground bud bank (root buds excluded): **19**
Depth of the belowground bud bank (root buds excluded) [cm]: **4**
Number of buds per shoot at the soil surface (root buds included): **4**
Number of buds per shoot at a depth of 0–10 cm (root buds included): **15**
Number of buds per shoot at a depth greater than 10 cm (root buds included): **0**
Size of the belowground bud bank (root buds included): **19**
Depth of the belowground bud bank (root buds included) [cm]: **4**

Trophic mode

Parasitism and mycoheterotrophy: **autotrophic**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Karyology

Chromosome number (2n): **30**

Ploidy level (x): **2**

2C genome size [Mbp]: **563.86**

1Cx monoploid genome size [Mbp]: **281.93**

Genomic GC content: **40.2 %**

Taxon origin

Origin in the Czech Republic: **native**

Ecological indicator values

Ellenberg-type indicator values

Light indicator value: **9 - full light plant, occurring only in fully irradiated places, not at less than 50% of diffuse radiation incident in an open area**

Temperature indicator value: **1 - cold indicator, occurring only in high mountain areas, i.e. the alpine and nival belts**

Moisture indicator value: **4 - transition between values 3 and 5**

Reaction indicator value: **2 - transition between values 1 and 3**

Nutrient indicator value: **2 - transition between values 1 and 3**

Salinity indicator value: **0 - not salt tolerant, glycophyte**

Habitat and sociology

Occurrence in habitats

1 Vegetation of cliffs, screes and walls

1B Siliceous cliffs and block fields: **1 - rare occurrence**

2 Alpine and subalpine grasslands

2A Alpine grasslands on siliceous bedrock: **2 - optimum**

11 Heathlands and scrub

11A Dry lowland to subalpine heathlands: **1 - rare occurrence**

Affinity to the forest environment

Affinity to the forest environment in Thermophyticum: **0 - taxon that does not spontaneously occur in Czech forests**

Affinity to the forest environment in Mesophyticum and Oreophyticum: **0 - taxon that does not spontaneously occur in Czech forests**

Diagnostic taxon

Diagnostic taxon of classes: [AA *Loiseleurio-Vaccinietea*](#)

Diagnostic taxon of alliances: [AAA *Loiseleurio procumbentis-Vaccinion*](#), [ABA *Juncion trifidi*](#)

Diagnostic taxon of associations: [AAA01 *Avenello flexuosae-Callunetum vulgaris*](#), [AAA02 *Junco trifidi-Empetretum hermaphroditi*](#), [ABA01 *Cetrario-Festucetum supinae*](#)

Ecological specialization indices

Ecological specialization index for all vegetation types: **7.1**

Ecological specialization index for non-forest vegetation: **7.1**

Distribution and frequency

Floristic zone: **arctic, boreal, northern temperate, southern temperate, submeridional**

Floristic region: **Europe, Western Siberia, Eastern America**

Distribution range extension along the continentality gradient: **5**

Elevational belt in the Czech Republic: **subalpine belt**

Occurrence frequency in the basic grid mapping cells and quadrants of the basic grid mapping cells: **13**

taxon.data.freq_in_quad: **16**

Commonness in vegetation plots from the Czech Republic

Occurrence frequency in vegetation plots: **0 %**

Occurrence frequency in vegetation plots with a cover above 5%: **42.9 %**

Occurrence frequency in vegetation plots with a cover above 25%: **0 %**

Occurrence frequency in vegetation plots with a cover above 50%: **0 %**

Mean percentage cover in vegetation plots: **7.4 %**

Maximum percentage cover in vegetation plots: **13 %**

Number of habitats with taxon occurrence in the Czech Republic

Number of narrow habitats in which the taxon occurs: **3**

Number of narrow habitats in which the taxon has its optimum: **1**

Number of broad habitats in which the taxon occurs: **3**

Number of broad habitats in which the taxon has its optimum: **1**

Threats and protection

Red List 2017 (national categories): **C2b - endangered taxon, rare and declining**

Red List 2017 (IUCN categories): **VU - vulnerable**

Legal protection: **not protected by law**