
Lepidium africanum

Distribution



Habitus and growth type

Life form: **therophyte**

Leaf

Leaf presence and metamorphosis: **leaves present, not modified**
Leaf arrangement (phyllotaxis): **alternate**
Leaf shape: **simple – entire**
Stipules: **absent**
Petiole: **absent**

Flower

Flower colour: **white**
Perianth type: **calyx present, corolla reduced or absent**
Perianth fusion: **reduced**
Calyx fusion: **aposepalous**
Inflorescence type: **racemus**
Generative reproduction type: **autogamy**

Fruit, seed and dispersal

Fruit type: **dry fruit – silicula**
Myrmecochory: **non-myrmecochorous (b)**

© 2014-2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Trophic mode

Parasitism and mycoheterotrophy: **autotrophic**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in the Czech Republic: **neophyte**
Invasion status: **casual**
Geographic origin: **Africa**
Year of the first record in the wild: **1964**
Year of the last record: **1990**
Period of introduction: **Recent Past (1950–2000)**
Introduction pathway: **unintentional – industry, unintentional – anthropogenic**

Habitat and sociology

Affinity to the forest environment

Affinity to the forest environment in Thermophyticum: **0 – taxon that does not spontaneously occur in Czech forests**
Affinity to the forest environment in Mesophyticum and Oreophyticum: **0 – taxon that does not spontaneously occur in Czech forests**

Threats and protection

Red List 2017 (national categories): **taxon is not on the Red List**
Red List 2017 (IUCN categories): **LC(NA) – least concern (taxon is not on the Red List)**
Legal protection: **not protected by law**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz