
Lepidium ruderale

Distribution





Map info

revised records

unrevised records

On the map are not visualized records without the coordinates and records marked as incorrect or doubtful.









Habitus and growth type

Height [m]: **0.1–0.5**
Growth form: **annual herb**
Life form: **therophyte**
Life strategy: **R – ruderal**
Life strategy (Pierce method based on leaf traits): **R**
Life strategy (Pierce method, C-score): **10.8 %**
Life strategy (Pierce method, S-score): **0 %**
Life strategy (Pierce method, R-score): **89.2 %**

Leaf

Leaf presence and metamorphosis: **leaves present, not modified**
Leaf arrangement (phyllotaxis): **alternate**
Leaf shape: **simple – entire, simple – pinnately divided**
Stipules: **absent**
Petiole: **both present and absent**
Leaf life span: **overwintering green**
Leaf anatomy: **scleromorphic, mesomorphic**

Flower

Flowering period [month]: **May–August**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz



Flowering phase: **6 Cornus sanguinea-Melica uniflora (start of early summer)**
Flower colour: **white**
Perianth type: **calyx present, corolla absent**
Perianth fusion: **reduced**
Calyx fusion: **aposepalous**
Inflorescence type: **racemus**
Dicliny: **synoecious**
Generative reproduction type: **autogamy**
Pollination syndrome: **selfing**

Fruit, seed and dispersal

Fruit type: **dry fruit – silicula**
Fruit colour: **brown**
Reproduction type: **only by seed/spores**
Dispersal unit (diaspore): **seed, fruit, infrutescence or its part**
Dispersal strategy: **Allium (mainly autochory)**
Myrmecochory: **non-myrmecochorous (b)**

Belowground organs and clonality

Shoot life span (cyclicity): **monocyclic shoots prevailing**
Primary root: **present**

Bud bank

Number of buds per shoot at the soil surface (root buds excluded): **5**
Number of buds per shoot at a depth of 0–10 cm (root buds excluded): **3**
Number of buds per shoot at a depth greater than 10 cm (root buds excluded): **0**
Size of the belowground bud bank (root buds excluded): **8**
Depth of the belowground bud bank (root buds exluded) [cm]: **2**
Number of buds per shoot at the soil surface (root buds included): **5**
Number of buds per shoot at a depth of 0–10 cm (root buds included): **3**
Number of buds per shoot at a depth greater than 10 cm (root buds included): **0**
Size of the belowground bud bank (root buds included): **8**
Depth of the belowground bud bank (root buds included) [cm]: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotrophic**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Karyology

Chromosome number (2n): **32 (16)**
Ploidy level (x): **4 (2)**
2C genome size [Mbp]: **586.57**
1Cx monoploid genome size [Mbp]: **146.64**
Genomic GC content: **37.1 %**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Taxon origin

Origin in the Czech Republic: **archaeophyte**
Invasion status: **naturalized**
Geographic origin: **Mediterranean**
Period of introduction: **Bronze Age (2300–750 BCE)**
Introduction pathway: **unintentional – anthropogenic**

Ecological indicator values

Ellenberg-type indicator values

Light indicator value: **9 – full light plant, occurring only in fully irradiated places, not at less than 50% of diffuse radiation incident in an open area**
Temperature indicator value: **6 – transition between values 5 and 7**
Moisture indicator value: **4 – transition between values 3 and 5**
Reaction indicator value: **7x – indicator of slightly acidic to slightly basic conditions, never occurring in very acidic conditions (generalist)**
Nutrient indicator value: **6 – transition between values 5 and 7**
Salinity indicator value: **1 – salt tolerant, mostly on low-salt to salt-free soils, but occasionally on slightly salty soils**

Indicator values for disturbance

Whole-community disturbance frequency indicator value: **0.29**
Herb layer disturbance frequency indicator value: **0.29**
Whole-community disturbance severity indicator value: **0.84**
Herb layer disturbance severity indicator value: **0.84**
Whole-community structure based disturbance indicator value: **0.67**
Herb layer structure-based disturbance indicator value: **0.67**

Habitat and sociology

Occurrence in habitats

6 Meadows and mesic pastures

6C Pastures and park grasslands: **2 – optimum**
6G Vegetation of wet disturbed soils: **1 – rare occurrence**

9 Sand grasslands and rock-outcrop vegetation

9B Open vegetation of acidic sands: **1 – rare occurrence**
9E Acidophilous vegetation of spring therophytes and succulents: **1 – rare occurrence**
9F Basiphilous vegetation of spring therophytes and succulents: **1 – rare occurrence**

13 Anthropogenic vegetation

13A Annual vegetation of ruderal habitats: **2 – optimum**
13C Annual vegetation of trampled habitats: **2 – optimum**
13D Perennial thermophilous ruderal vegetation: **1 – rare occurrence**

Affinity to the forest environment

Affinity to the forest environment in Thermophyticum: **0 – taxon that does not spontaneously occur in Czech forests**
Affinity to the forest environment in Mesophyticum and Oreophyticum: **0 – taxon that does not spontaneously occur in Czech forests**

Diagnostic taxon

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Diagnostic taxon of classes: **XA *Polygono arenastri-Poëtea annuae***
Diagnostic taxon of alliances: **XAA *Coronopodo-Polygonion arenastri*, XBH *Sisymbrion officinalis*, XBK *Eragrostion cilianensi-minoris***
Diagnostic taxon of associations: **XAA02 *Sclerochloo durae-Polygonetum arenastri*, XAA04 *Eragrostio minoris-Polygonetum arenastri*, XBG08 *Descurainietum sophiae*, XBG12 *Ivaetum xanthiifoliae*, XBH01 *Hordeetum murini*, XBK01 *Digitario sanguinalis-Eragrostietum minoris*, XBK03 *Eragrostio poaeoidis-Panicetum capillaris***

Constant taxon

Constant taxon of associations: **XAA02 *Sclerochloo durae-Polygonetum arenastri*, XBI03 *Polygono arenastri-Chenopodietum muralis*, XBK01 *Digitario sanguinalis-Eragrostietum minoris*, XBK03 *Eragrostio poaeoidis-Panicetum capillaris***

Dominant taxon

Dominant taxon of associations: **XBG02 *Chenopodietum urbici***

Ecological specialization indices

Ecological specialization index for all vegetation types: **5.1**
Ecological specialization index for non-forest vegetation: **5.1**

Distribution and frequency

Floristic zone: **boreal, northern temperate, southern temperate, submeridional, meridional**
Floristic region: **Europe, Asia**
Continentality degree: **6**
Distribution range extension along the continentality gradient: **6**
Elevational belt in the Czech Republic: **lowlands, colline belt (montane belt)**
Occurrence frequency in the basic grid mapping cells and quadrants of the basic grid mapping cells: 509
taxon.data.freq_in_quad: 1162

Commonness in vegetation plots from the Czech Republic

Occurrence frequency in vegetation plots: **0.7 %**
Occurrence frequency in vegetation plots with a cover above 5%: **5.1 %**
Occurrence frequency in vegetation plots with a cover above 25%: **1.9 %**
Occurrence frequency in vegetation plots with a cover above 50%: **0.9 %**
Mean percentage cover in vegetation plots: **3.5 %**
Maximum percentage cover in vegetation plots: **63 %**

Number of habitats with taxon occurrence in the Czech Republic

Number of narrow habitats in which the taxon occurs: **8**
Number of narrow habitats in which the taxon has its optimum: **3**
Number of broad habitats in which the taxon occurs: **3**
Number of broad habitats in which the taxon has its optimum: **2**

Threats and protection

Red List 2017 (national categories): **taxon is not on the Red List**
Red List 2017 (IUCN categories): **LC(NA) – least concern (taxon is not on the Red List)**
Legal protection: **not protected by law**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz