
Lysimachia nemorum

Distribution





Map info

revised records

unrevised records

On the map are not visualized records without the coordinates and records marked as incorrect or doubtful.







Habitus and growth type

Height [m]: **0.1–0.4**
Growth form: **clonal herb**
Life form: **hemicryptophyte**
Life strategy: **CS – competitor/stress-tolerator**
Life strategy (Pierce method based on leaf traits): **R/SR**
Life strategy (Pierce method, C-score): **14.3 %**
Life strategy (Pierce method, S-score): **20.7 %**
Life strategy (Pierce method, R-score): **65 %**

Leaf

Leaf presence and metamorphosis: **leaves present, not modified**
Leaf arrangement (phyllotaxis): **opposite**
Leaf shape: **simple – entire**
Stipules: **absent**
Petiole: **present**
Leaf life span: **evergreen**
Leaf anatomy: **hygromorphic, helomorphic**

Flower

Flowering period [month]: **June–August**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz



Flowering phase: **5 Sorbus aucuparia-Galium odoratum (end of mid-spring)**
Flower colour: **yellow**
Flower symmetry: **actinomorphic**
Perianth type: **calyx and corolla**
Perianth fusion: **fused**
Shape of the sympetalous corolla or syntepalous perianth: **rotate, funnel-shaped**
Calyx fusion: **fused at the base**
Inflorescence type: **flores solitarii**
Dicliny: **synoecious**
Pollination syndrome: **insect-pollination, selfing**



Fruit, seed and dispersal

Fruit type: **dry fruit – capsule**
Fruit colour: **brown**
Reproduction type: **by seed/spores and vegetatively**
Dispersal unit (diaspore): **seed**
Dispersal strategy: **Allium (mainly autochory)**
Myrmecochory: **non-myrmecochorous (b)**

Belowground organs and clonality

Shoot metamorphosis: **stolon**
Storage organ: **stolon**
Type of clonal growth organ: **stolon**
Freely dispersible organs of clonal growth: **absent**
Shoot life span (cyclicity): **monocyclic shoots prevailing**
Branching type of stem-derived organs of clonal growth: **monopodial**
Primary root: **absent**
Persistence of the clonal growth organ [year]: **2**
Number of clonal offspring: **4.2**
Lateral spreading distance by clonal growth [m]: **0.13**
Clonal index: **5**

Bud bank

Number of buds per shoot at the soil surface (root buds excluded): **15**
Number of buds per shoot at a depth of 0–10 cm (root buds excluded): **0**
Number of buds per shoot at a depth greater than 10 cm (root buds excluded): **0**
Size of the belowground bud bank (root buds excluded): **15**
Depth of the belowground bud bank (root buds exluded) [cm]: **1**
Number of buds per shoot at the soil surface (root buds included): **15**
Number of buds per shoot at a depth of 0–10 cm (root buds included): **0**
Number of buds per shoot at a depth greater than 10 cm (root buds included): **0**
Size of the belowground bud bank (root buds included): **15**
Depth of the belowground bud bank (root buds included) [cm]: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotrophic**
Carnivory: **non-carnivorous**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in the Czech Republic: **native**

Ecological indicator values

Ellenberg-type indicator values

Light indicator value: **2 – transition between values 1 and 3**
Temperature indicator value: **4 – transition between values 3 and 5**
Moisture indicator value: **7 – humidity indicator, focus on well moistened, but not wet soils**
Reaction indicator value: **5 – indicator of moderate acidity, occurring rarely in strongly acidic as well as in neutral to alkaline conditions**
Nutrient indicator value: **6 – transition between values 5 and 7**
Salinity indicator value: **0 – not salt tolerant, glycophyte**

Indicator values for disturbance

Whole-community disturbance frequency indicator value: **-1.71**
Herb layer disturbance frequency indicator value: **-0.6**
Whole-community disturbance severity indicator value: **0.28**
Herb layer disturbance severity indicator value: **0.21**
Whole-community structure based disturbance indicator value: **0.24**
Herb layer structure-based disturbance indicator value: **0.28**

Habitat and sociology

Occurrence in habitats

2 Alpine and subalpine grasslands

2B Subalpine tall-forb and tall-grass vegetation: **1 – rare occurrence**

4 Wetland and riverine herbaceous vegetation

4E Reed vegetation of brooks: **1 – rare occurrence**
4K Petasites fringes of montane brooks: **2 – optimum**

5 Vegetation of springs and mires

5B Lowland to montane soft-water springs: **2 – optimum**
5C Alpine and subalpine soft-water springs: **1 – rare occurrence**
5D Calcareous fens: **1 – rare occurrence**
5E Acidic moss-rich fens and peatland meadows: **1 – rare occurrence**
5F Transitional mires: **1 – rare occurrence**

6 Meadows and mesic pastures

6B Montane mesic meadows: **1 – rare occurrence**
6E Wet Cirsium meadows: **1 – rare occurrence**
6G Vegetation of wet disturbed soils: **2 – optimum**

7 Acidophilous grasslands

7A Subalpine and montane acidophilous grasslands: **1 – rare occurrence**
7B Submontane Nardus grasslands: **1 – rare occurrence**

11 Heathlands and scrub

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

11H Subalpine deciduous scrub: **1 – rare occurrence**
11R Scrub and pioneer woodland of forests clearings: **1 – rare occurrence**

12 Forests

12B Alluvial forests: **2 – optimum**
12D Ravine forests: **1 – rare occurrence**
12E Herb-rich beech forests: **2 – optimum**
12G Acidophilous beech forests: **1 – rare occurrence**
12R Acidophilous spruce forests: **1 – rare occurrence**
12S Basiphilous spruce forests: **1 – rare occurrence**
12U Plantations of broad-leaved non-native trees: **1 – rare occurrence**
12V Spruce plantations: **1 – rare occurrence**

13 Anthropogenic vegetation

13E Perennial nitrophilous herbaceous vegetation of mesic sites: **1 – rare occurrence**
13F Herbaceous vegetation of forests clearings and Rubus scrub: **1 – rare occurrence**

Affinity to the forest environment

Affinity to the forest environment in Mesophyticum and Oreophyticum: **1.1 – taxon occurring mainly in the closed forest**

Diagnostic taxon

Diagnostic taxon of classes: **RA *Montio-Cardaminetea***
Diagnostic taxon of alliances: **RAA *Caricion remotae***
Diagnostic taxon of associations: **LBA02 *Piceo abietis-Alnetum glutinosae***, **LBA03 *Carici remotae-Fraxinetum excelsioris***, **LBC04 *Athyrio distentifolii-Fagetum sylvaticae***

Constant taxon

Constant taxon of associations: **LBC04 *Athyrio distentifolii-Fagetum sylvaticae***

Ecological specialization indices

Ecological specialization index for all vegetation types: **5.1**
Ecological specialization index for non-forest vegetation: **4.7**
Ecological specialization index for forest vegetation: **5.9**

Colonization ability

Index of colonization success (ICS): **4**
Index of colonization potential (ICP): **3**
Optimum successional age [years]: **5**

Distribution and frequency

Floristic zone: **northern temperate, southern temperate, submeridional, meridional**
Floristic region: **Europe**
Continentality degree: **3**
Distribution range extension along the continentality gradient: **3**
Elevational belt in the Czech Republic: **submontane belt, montane belt**
Occurrence frequency in the basic grid mapping cells and quadrants of the basic grid mapping cells: 541
taxon.data.freq_in_quad: 1473

Commonness in vegetation plots from the Czech Republic

Occurrence frequency in vegetation plots: **1.5 %**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Occurrence frequency in vegetation plots with a cover above 5%: **9.4 %**
Occurrence frequency in vegetation plots with a cover above 25%: **1.4 %**
Occurrence frequency in vegetation plots with a cover above 50%: **0.5 %**
Mean percentage cover in vegetation plots: **3.6 %**
Maximum percentage cover in vegetation plots: **88 %**

Number of habitats with taxon occurrence in the Czech Republic

Number of narrow habitats in which the taxon occurs: **25**
Number of narrow habitats in which the taxon has its optimum: **5**
Number of broad habitats in which the taxon occurs: **8**
Number of broad habitats in which the taxon has its optimum: **4**

Threats and protection

Red List 2017 (national categories): **taxon is not on the Red List**
Red List 2017 (IUCN categories): **LC(NA) – least concern (taxon is not on the Red List)**
Legal protection: **not protected by law**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz