
# *Oxalis debilis*

## Distribution




### Habitus and growth type

Growth form: **clonal herb**
Life form: **geophyte (hemicryptophyte)**

### Leaf

Leaf presence and metamorphosis: **leaves present, not modified**
Leaf arrangement (phyllotaxis): **rosulate**
Leaf shape: **compound – ternate**
Stipules: **absent**
Petiole: **present**

### Flower

Flower colour: **pink-violet, blue-violet**
Flower symmetry: **actinomorphic**
Perianth type: **calyx and corolla**
Perianth fusion: **free, fused**
Shape of the sympetalous corolla or syntepalous perianth: **funnel-shaped**
Calyx fusion: **aposepalous**
Inflorescence type: **pseudumbella**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*

## Fruit, seed and dispersal

Fruit type: **dry fruit – capsule**
Dispersal strategy: **Zea (no dispersal)**
Myrmecochory: **probably myrmecochorous nv**

## Belowground organs and clonality

Type of clonal growth organ: **bulb**
Freely dispersible organs of clonal growth: **absent**
Shoot life span (cyclicity): **monocyclic shoots prevailing**
Branching type of stem-derived organs of clonal growth: **sympodial**
Primary root: **absent**
Number of clonal offspring: **3.3**
Lateral spreading distance by clonal growth [m]: **0.01**
Clonal index: **4**

### Bud bank

Number of buds per shoot at the soil surface (root buds excluded): **5**
Number of buds per shoot at a depth of 0–10 cm (root buds excluded): **15**
Number of buds per shoot at a depth greater than 10 cm (root buds excluded): **0**
Size of the belowground bud bank (root buds excluded): **20**
Depth of the belowground bud bank (root buds exluded) [cm]: **4**
Number of buds per shoot at the soil surface (root buds included): **5**
Number of buds per shoot at a depth of 0–10 cm (root buds included): **15**
Number of buds per shoot at a depth greater than 10 cm (root buds included): **0**
Size of the belowground bud bank (root buds included): **20**
Depth of the belowground bud bank (root buds included) [cm]: **4**

## Trophic mode

Parasitism and mycoheterotrophy: **autotrophic**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in the Czech Republic: **neophyte**
Invasion status: **casual**
Geographic origin: **South America**
Year of the first record in the wild: **1967**
Year of the last record: **1979**
Period of introduction: **Recent Past (1950–2000)**
Introduction pathway: **intentional – ornamental, unintentional – horticulture**

## Habitat and sociology

### Affinity to the forest environment

Affinity to the forest environment in Thermophyticum: **0 – taxon that does not spontaneously occur in Czech forests**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*

Affinity to the forest environment in Mesophyticum and Oreophyticum: **0 – taxon that does not spontaneously occur in Czech forests**

## Threats and protection

Red List 2017 (national categories): **taxon is not on the Red List**
Red List 2017 (IUCN categories): **LC(NA) – least concern (taxon is not on the Red List)**
Legal protection: **not protected by law**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*