


Rumex arifolius

Distribution



© Pavel Veselý

Map info

● revised records

○ unrevised records

On the map are not visualized records without the coordinates and records marked as incorrect or doubtful.

Habitus and growth type

Height [m]: **0.4-1**Growth form: **clonal herb**Life form: **hemicryptophyte**Life strategy: **C - competitor**Life strategy (Pierce method based on leaf traits): **CR**Life strategy (Pierce method, C-score): **56.8 %**Life strategy (Pierce method, S-score): **0 %**Life strategy (Pierce method, R-score): **43.2 %**

Leaf

Leaf presence and metamorphosis: **leaves present, not modified**Leaf arrangement (phyllotaxis): **alternate**Leaf shape: **simple - entire**Stipules: **present**Petiole: **both present and absent**Leaf life span: **summer green**Leaf anatomy: **succulent, mesomorphic**

© Eva Hlubíková



© Pavel Veselý

Flower

Flowering period [month]: **June-September**

Flowering phase: **6 Cornus sanguinea-Melica uniflora (start of early summer)**

Flower colour: **green**

Flower symmetry: **actinomorphic**

Perianth type: **homochlamydeous**

Perianth fusion: **free**

Inflorescence type: **panicula e pseudospicis composita**

Dicliny: **dioecious**

Generative reproduction type: **alogamy**

Pollination syndrome: **wind-pollination**

Fruit, seed and dispersal

Fruit type: **dry fruit - achene/cypsela/samara**

Fruit colour: **yellow, red, brown, grey**

Reproduction type: **only by seed/spores**

Dispersal unit (diaspore): **fruit, infructescence or its part**

Dispersal strategy: **Allium (mainly autochory)**

Myrmecochory: **non-myrmecochorous (b)**

Belowground organs and clonality

Shoot metamorphosis: **pleiocorm**

Root metamorphosis: **primary storage root**

Storage organ: **pleiocorm, primary storage root**

Type of clonal growth organ: **epigeogenous rhizome**

Freely dispersible organs of clonal growth: **absent**

Shoot life span (cyclicality): **dicyclic or polycyclic shoots prevailing**

Branching type of stem-derived organs of clonal growth: **sympodial**

Primary root: **absent**

Persistence of the clonal growth organ [year]: **4**

Number of clonal offspring: **2.1**

Lateral spreading distance by clonal growth [m]: **0.01**

Clonal index: **3**

Bud bank

Number of buds per shoot at the soil surface (root buds excluded): **5**

Number of buds per shoot at a depth of 0–10 cm (root buds excluded): **15**

Number of buds per shoot at a depth greater than 10 cm (root buds excluded): **0**

Size of the belowground bud bank (root buds excluded): **20**

Depth of the belowground bud bank (root buds excluded) [cm]: **4**

Number of buds per shoot at the soil surface (root buds included): **5**

Number of buds per shoot at a depth of 0–10 cm (root buds included): **15**

Number of buds per shoot at a depth greater than 10 cm (root buds included): **0**

Size of the belowground bud bank (root buds included): **20**

Depth of the belowground bud bank (root buds included) [cm]: **4**

Trophic mode

Parasitism and mycoheterotrophy: **autotrophic**

Carnivory: **non-carnivorous**



Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Karyology

Chromosome number (2n): **14**

Ploidy level (x): **2**

2C genome size [Mbp]: **5911.89**

1Cx monoploid genome size [Mbp]: **2955.95**

Genomic GC content: **44.1 %**

Taxon origin

Origin in the Czech Republic: **native**

Ecological indicator values

Ellenberg-type indicator values

Light indicator value: **6 - transition between values 5 and 7; rarely at less than 20% of diffuse radiation incident in an open area**

Temperature indicator value: **3 - cool indicator, occurring mainly in subalpine areas**

Moisture indicator value: **6 - transition between values 5 and 7**

Reaction indicator value: **6 - transition between values 5 and 7**

Nutrient indicator value: **6 - transition between values 5 and 7**

Salinity indicator value: **0 - not salt tolerant, glycophyte**

Indicator values for disturbance

Whole-community disturbance frequency indicator value: **-1.44**

Herb layer disturbance frequency indicator value: **-0.68**

Whole-community disturbance severity indicator value: **0.22**

Herb layer disturbance severity indicator value: **0.21**

Whole-community structure based disturbance indicator value: **0.34**

Herb layer structure-based disturbance indicator value: **0.39**

Habitat and sociology

Occurrence in habitats

2 Alpine and subalpine grasslands

2A Alpine grasslands on siliceous bedrock: **1 - rare occurrence**

2B Subalpine tall-forb and tall-grass vegetation: **2 - optimum**

4 Wetland and riverine herbaceous vegetation

4K Petasites fringes of montane brooks: **2 - optimum**

5 Vegetation of springs and mires

5B Lowland to montane soft-water springs: **1 - rare occurrence**

5C Alpine and subalpine soft-water springs: **2 - optimum**

5E Acidic moss-rich fens and peatland meadows: **1 - rare occurrence**

6 Meadows and mesic pastures

6B Montane mesic meadows: **2 - optimum**

7 Acidophilous grasslands

7A Subalpine and montane acidophilous grasslands: **2 - optimum**

7B Submontane Nardus grasslands: **1 - rare occurrence**

11 Heathlands and scrub

11A Dry lowland to subalpine heathlands: **1 - rare occurrence**

11D Subalpine acidophilous Pinus mugo scrub: **1 - rare occurrence**

11H Subalpine deciduous scrub: **2 - optimum**

12 Forests

12B Alluvial forests: **1 - rare occurrence**

12E Herb-rich beech forests: **1 - rare occurrence**

12G Acidophilous beech forests: **1 - rare occurrence**

12R Acidophilous spruce forests: **1 - rare occurrence**

12S Basiphilous spruce forests: **2 - optimum**

13 Anthropogenic vegetation

13E Perennial nitrophilous herbaceous vegetation of mesic sites: **1 - rare occurrence**

Affinity to the forest environment

Affinity to the forest environment in Mesophyticum and Oreophyticum: **2.1 - taxon occurring both in the forest and open vegetation**

Diagnostic taxon

Diagnostic taxon of classes: [AD Mulgedio-Aconitetea](#)

Diagnostic taxon of alliances: [ADA Calamagrostion villosae](#), [ADB Calamagrostion arundinaceae](#), [ADC Salicion silesiaca](#), [ADD Adenostylion alliariae](#), [ADE Dryopterido filicis-maris-Athyrium distentifolii](#), [RAD Swertio perennis-Dichodontion palustris](#), [XDF Rumicion alpini](#)

Diagnostic taxon of associations: [ADA02 Crepido conyzifoliae-Calamagrostietum villosae](#), [ADA03 Viola sudeticae-Deschampsietum cespitosae](#), [ADB01 Bupleuro longifoliae-Calamagrostietum arundinaceae](#), [ADC01 Salici silesiaca-Betuletum carpatica](#), [ADC02 Pado borealis-Sorbetum aucupariae](#), [ADD01 Ranunculo platanifolii-Adenostyletum alliariae](#), [ADD02 Salicetum lapponum](#), [ADD03 Trollio altissimi-Geranium sylvatici](#), [ADD04 Laserpitio archangelicae-Dactylidetum glomeratae](#), [ADE01 Daphno mezerei-Dryopteridetum filicis-maris](#), [ADE02 Adenostylo alliariae-Athyrium distentifolii](#), [LBA01 Alnetum incanae](#), [LBC04 Athyrio distentifolii-Fagetum sylvaticae](#), [LFC02 Athyrio distentifolii-Piceetum abietis](#), [RAD03 Cardaminetum opicii](#), [XDF01 Rumicetum alpini](#)

Constant taxon

Constant taxon of classes: [AD Mulgedio-Aconitetea](#)

Constant taxon of alliances: [ADA Calamagrostion villosae](#), [ADC Salicion silesiaca](#), [ADD Adenostylion alliariae](#), [ADE Dryopterido filicis-maris-Athyrium distentifolii](#), [XDF Rumicion alpini](#)

Constant taxon of associations: [ADA02 Crepido conyzifoliae-Calamagrostietum villosae](#), [ADA03 Viola sudeticae-Deschampsietum cespitosae](#), [ADC01 Salici silesiaca-Betuletum carpatica](#), [ADC02 Pado borealis-Sorbetum aucupariae](#), [ADD01 Ranunculo platanifolii-Adenostyletum alliariae](#), [ADD02 Salicetum lapponum](#), [ADD04 Laserpitio archangelicae-Dactylidetum glomeratae](#), [ADE01 Daphno mezerei-Dryopteridetum filicis-maris](#), [ADE02 Adenostylo alliariae-Athyrium distentifolii](#), [KCA02 Adenostylo alliariae-Pinetum mugo](#), [LBC04 Athyrio distentifolii-Fagetum sylvaticae](#), [LFC02 Athyrio distentifolii-Piceetum abietis](#), [XDF01 Rumicetum alpini](#)

Ecological specialization indices

Ecological specialization index for all vegetation types: **4.6**

Ecological specialization index for non-forest vegetation: **4.5**

Ecological specialization index for forest vegetation: **6**

Distribution and frequency

Floristic zone: **arctic, boreal, northern temperate, southern temperate, submeridional**

Floristic region: **Europe, Asia**

Continentality degree: **6**

Distribution range extension along the continentality gradient: **5**

Elevational belt in the Czech Republic: **montane belt, subalpine belt**

Occurrence frequency in the basic grid mapping cells and quadrants of the basic grid mapping cells: **89**

taxon.data.freq_in_quad: **184**

Commonness in vegetation plots from the Czech Republic

Occurrence frequency in vegetation plots: **0.7 %**

Occurrence frequency in vegetation plots with a cover above 5%: **10.4 %**

Occurrence frequency in vegetation plots with a cover above 25%: **1.4 %**

Occurrence frequency in vegetation plots with a cover above 50%: **0 %**

Mean percentage cover in vegetation plots: **3.8 %**

Maximum percentage cover in vegetation plots: **42 %**

Number of habitats with taxon occurrence in the Czech Republic

Number of narrow habitats in which the taxon occurs: **18**

Number of narrow habitats in which the taxon has its optimum: **7**

Number of broad habitats in which the taxon occurs: **8**

Number of broad habitats in which the taxon has its optimum: **7**

Threats and protection

Red List 2017 (national categories): **taxon is not on the Red List**

Red List 2017 (IUCN categories): **LC(NA) - least concern (taxon is not on the Red List)**

Legal protection: **not protected by law**