
Filipendula ulmaria subsp. *ulmaria*

Distribution



Map info

revised records

unrevised records

On the map are not visualized records without the coordinates and records marked as incorrect or doubtful.

Leaf

Leaf presence and metamorphosis: **leaves present, not modified**
Leaf arrangement (phyllotaxis): **alternate**
Leaf shape: **compound – interruptedly pinnate**
Stipules: **present**
Petiole: **present**

Flower

Flower colour: **yellow-white**
Flower symmetry: **actinomorphic**
Perianth type: **calyx and corolla**
Perianth fusion: **free**
Calyx fusion: **aposepalous**
Inflorescence type: **anthella**

Fruit, seed and dispersal

Fruit type: **dry fruit – head of achenes**
Dispersal unit (diaspore): **fruit, infrutescence or its part**
Dispersal strategy: **Allium (mainly autochory)**
Myrmecochory: **non-myrmecochorous (b)**

© 2014-2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Trophic mode

Parasitism and mycoheterotrophy: **autotrophic**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in the Czech Republic: **native**

Ecological indicator values

Ellenberg-type indicator values

Light indicator value: **6 – transition between values 5 and 7; rarely at less than 20% of diffuse radiation incident in an open area**
Temperature indicator value: **5x – moderate heat indicator, occurring from lowland to montane belt, mainly in submontane-temperate areas (generalist)**
Moisture indicator value: **8 – transition between values 7 and 9**
Reaction indicator value: **6x – transition between values 5 and 7 (generalist)**
Nutrient indicator value: **5 – occurring at moderately nutrient-rich sites, and less frequently at poor and rich sites**
Salinity indicator value: **1 – salt tolerant, mostly on low-salt to salt-free soils, but occasionally on slightly salty soils**

Habitat and sociology

Occurrence in habitats

2 Alpine and subalpine grasslands

2B Subalpine tall-forb and tall-grass vegetation: **1 – rare occurrence**

4 Wetland and riverine herbaceous vegetation

4A Reed-beds of eutrophic still waters: **1 – rare occurrence**
4B Halophilous reed and sedge beds: **1 – rare occurrence**
4D Riverine reed vegetation: **2 – optimum**
4E Reed vegetation of brooks: **1 – rare occurrence**
4F Mesotrophic vegetation of muddy substrata: **1 – rare occurrence**
4G Tall-sedge beds: **2 – optimum**
4J River gravel banks: **1 – rare occurrence**
4K Petasites fringes of montane brooks: **2 – optimum**
4L Nitrophilous herbaceous fringes of lowland rivers: **1 – rare occurrence**

5 Vegetation of springs and mires

5B Lowland to montane soft-water springs: **1 – rare occurrence**
5C Alpine and subalpine soft-water springs: **1 – rare occurrence**
5D Calcareous fens: **2 – optimum**
5E Acidic moss-rich fens and peatland meadows: **2 – optimum**
5F Transitional mires: **1 – rare occurrence**

6 Meadows and mesic pastures

6A Mesic Arrhenatherum meadows: **1 – rare occurrence**
6B Montane mesic meadows: **1 – rare occurrence**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

6C Pastures and park grasslands: **1 – rare occurrence**
6D Alluvial meadows of lowland rivers: **2 – optimum**
6E Wet Cirsium meadows: **3 – dominant**
6F Intermittently wet Molinia meadows: **2 – optimum**

7 Acidophilous grasslands

7B Submontane Nardus grasslands: **1 – rare occurrence**

9 Sand grasslands and rock-outcrop vegetation

9E Acidophilous vegetation of spring therophytes and succulents: **1 – rare occurrence**

10 Saline vegetation

10J Saline steppes: **1 – rare occurrence**

11 Heathlands and scrub

11H Subalpine deciduous scrub: **1 – rare occurrence**
11I Willow carrs: **1 – rare occurrence**
11J Willow galleries of loamy and sandy river banks: **2 – optimum**
11L Tall mesic and xeric shrub: **1 – rare occurrence**
11R Scrub and pioneer woodland of forests clearings: **1 – rare occurrence**

12 Forests

12A Alder carrs: **2 – optimum**
12B Alluvial forests: **2 – optimum**
12U Plantations of broad-leaved non-native trees: **1 – rare occurrence**

13 Anthropogenic vegetation

13E Perennial nitrophilous herbaceous vegetation of mesic sites: **1 – rare occurrence**

Affinity to the forest environment

Affinity to the forest environment in Thermophyticum: **2.1 – taxon occurring both in the forest and open vegetation**
Affinity to the forest environment in Mesophyticum and Oreophyticum: **2.1 – taxon occurring both in the forest and open vegetation**

Distribution and frequency

Elevational belt in the Czech Republic: **lowlands, colline belt, submontane belt, montane belt**
Occurrence frequency in the basic grid mapping cells and quadrants of the basic grid mapping cells: 461
taxon.data.freq_in_quad: 965

Number of habitats with taxon occurrence in the Czech Republic

Number of narrow habitats in which the taxon occurs: **33**
Number of narrow habitats in which the taxon has its optimum: **11**
Number of broad habitats in which the taxon occurs: **10**
Number of broad habitats in which the taxon has its optimum: **5**

Threats and protection

Red List 2017 (national categories): **taxon is not on the Red List**
Red List 2017 (IUCN categories): **LC(NA) – least concern (taxon is not on the Red List)**
Legal protection: **not protected by law**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz