
Odontites vernus subsp. *vernus*

Distribution





Map info

revised records

unrevised records

On the map are not visualized records without the coordinates and records marked as incorrect or doubtful.





Habitus and growth type

Height [m]: **0.1–0.4**
Life form: **therophyte**

Leaf

Leaf presence and metamorphosis: **leaves present, not modified**
Leaf arrangement (phyllotaxis): **opposite**
Leaf shape: **simple – entire**
Stipules: **absent**
Petiole: **absent**
Leaf life span: **summer green**
Leaf anatomy: **mesomorphic**

Flower

Flowering period [month]: **June–July**
Flower colour: **red-violet**
Flower symmetry: **zygomorphic**
Perianth type: **calyx and corolla**
Perianth fusion: **fused**
Shape of the sympetalous corolla or syntepalous perianth: **bilabiate**
Calyx fusion: **synsepalous**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Inflorescence type: **racemus**
Pollination syndrome: **insect-pollination, selfing**

Fruit, seed and dispersal

Fruit type: **dry fruit – capsule**
Reproduction type: **only by seed/spores**
Dispersal unit (diaspore): **seed**
Dispersal strategy: **Allium (mainly autochory)**
Myrmecochory: **myrmecochorous**

Trophic mode

Parasitism and mycoheterotrophy: **root hemiparasite**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in the Czech Republic: **native**

Ecological indicator values

Ellenberg-type indicator values

Light indicator value: **6 – transition between values 5 and 7; rarely at less than 20% of diffuse radiation incident in an open area**
Temperature indicator value: **6 – transition between values 5 and 7**
Moisture indicator value: **5 – indicator of fresh soils, focus on soils of average moisture, missing on wet and on soils that frequently dry out**
Reaction indicator value: **6 – transition between values 5 and 7**
Nutrient indicator value: **6 – transition between values 5 and 7**
Salinity indicator value: **1 – salt tolerant, mostly on low-salt to salt-free soils, but occasionally on slightly salty soils**

Habitat and sociology

Affinity to the forest environment

Affinity to the forest environment in Thermophyticum: **0 – taxon that does not spontaneously occur in Czech forests**
Affinity to the forest environment in Mesophyticum and Oreophyticum: **0 – taxon that does not spontaneously occur in Czech forests**

Distribution and frequency

Elevational belt in the Czech Republic: **lowlands, colline belt, submontane belt, montane belt**
Occurrence frequency in the basic grid mapping cells and quadrants of the basic grid mapping cells: 115
taxon.data.freq_in_quad: 167

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Threats and protection

Red List 2017 (national categories): **C2t – endangered taxon, declining**
Red List 2017 (IUCN categories): **EN – endangered**
Legal protection: **not protected by law**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz