Epilobium ×vicinum

Distribution



Habitus and growth type

Life form: **hemicryptophyte**

Leaf

Leaf presence and metamorphosis: **leaves present, not modified**
Stipules: **absent**
Petiole: **both present and absent**

Flower

Flower colour: **pink**
Perianth type: **calyx and corolla**
Calyx fusion: **hypanthium**
Inflorescence type: **racemus**

Trophic mode

Parasitism and mycoheterotrophy: **autotrophic**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Taxon origin

Origin in the Czech Republic: **neophyte**
Invasion status: **casual**
Geographic origin: **hybrid origin**
Year of the first record in the wild: **1971**
Period of introduction: **Early Modern Period (1500–1800)**
Introduction pathway: **unintentional – nature**

Habitat and sociology

Affinity to the forest environment

Affinity to the forest environment in Mesophyticum and Oreophyticum: **2.2 – taxon occurring partly in the forest, but mainly in open vegetation**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz