


# *Saxifraga* × *geum*

## Distribution



### Map info

● revised records

○ unrevised records

On the map are not visualized records without the coordinates and records marked as incorrect or doubtful.

## Habitus and growth type

Height [m]: **0.2-0.4**

Life form: **hemicryptophyte**

Life strategy: **CSR - competitor/stress-tolerator/ruderal**

## Leaf

Leaf presence and metamorphosis: **leaves present, not modified**

Stipules: **absent**

Petiole: **present**

Leaf life span: **evergreen**

## Flower

Flowering period [month]: **June-August**

Flower colour: **white**

Perianth type: **calyx and corolla**

Calyx fusion: **synsepalous**

Inflorescence type: **anthella**

Dicliny: **synoecious**

Pollination syndrome: **insect-pollination**

## **Fruit, seed and dispersal**

Reproduction type: **by seed/spores and vegetatively**

## **Belowground organs and clonality**

Shoot metamorphosis: **stolon**

Storage organ: **stolon**

## **Trophic mode**

Parasitism and mycoheterotrophy: **autotrophic**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## **Taxon origin**

Origin in the Czech Republic: **neophyte**

Invasion status: **casual**

Geographic origin: **hybrid origin**

Period of introduction: **Early Modern Period (1500-1800)**

Introduction pathway: **intentional - ornamental, intentional - nature**

## **Habitat and sociology**

Affinity to the forest environment

Affinity to the forest environment in Thermophyticum: **0 - taxon that does not spontaneously occur in Czech forests**

Affinity to the forest environment in Mesophyticum and Oreophyticum: **0 - taxon that does not spontaneously occur in Czech forests**

## **Distribution and frequency**

Floristic zone: **northern temperate, southern temperate, submeridional**

Floristic region: **Europe**

Occurrence frequency in the basic grid mapping cells and quadrants of the basic grid mapping cells: 3

taxon.data.freq\_in\_quad: 3