


Hordeum vulgare Distichon Group

Distribution



Map info

● revised records

○ unrevised records

On the map are not visualized records without the coordinates and records marked as incorrect or doubtful.

Habitus and growth type

Life form: **therophyte**

Life strategy (Pierce method based on leaf traits): **CR**

Life strategy (Pierce method, C-score): **44.1 %**

Life strategy (Pierce method, S-score): **5.6 %**

Life strategy (Pierce method, R-score): **50.3 %**

Leaf

Leaf presence and metamorphosis: **leaves present, not modified**

Petiole: **absent**

Flower

Flower colour: **green**

Perianth type: **reduced**

Inflorescence type: **spica e spiculis composita**

Fruit, seed and dispersal

Fruit colour: **brown**

Dispersal unit (diaspore): **fruit, infrutescence or its part**

Dispersal strategy: **Allium (mainly autochory)**

Trophic mode

Parasitism and mycoheterotrophy: **autotrophic**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in the Czech Republic: **archaeophyte**

Invasion status: **casual**

Geographic origin: **anecophyte**

Period of introduction: **Eneolithic (4200-2300 BCE)**

Introduction pathway: **intentional - crops**

Habitat and sociology

Affinity to the forest environment

Affinity to the forest environment in Thermophyticum: **0 - taxon that does not spontaneously occur in Czech forests**

Affinity to the forest environment in Mesophyticum and Oreophyticum: **0 - taxon that does not spontaneously occur in Czech forests**