Hordeum vulgare Distichon Group

Distribution



Map info

revised records

unrevised records

On the map are not visualized records without the coordinates and records marked as incorrect or doubtful.

Habitus and growth type

Life form: **therophyte**
Life strategy (Pierce method based on leaf traits): **CR**
Life strategy (Pierce method, C-score): **44.1 %**
Life strategy (Pierce method, S-score): **5.6 %**
Life strategy (Pierce method, R-score): **50.3 %**

Leaf

Leaf presence and metamorphosis: **leaves present, not modified**
Petiole: **absent**

Flower

Flower colour: **green**
Perianth type: **reduced**
Inflorescence type: **spica e spiculis composita**

Fruit, seed and dispersal

Fruit colour: **brown**
Dispersal unit (diaspore): **fruit, infrutescence or its part**
Dispersal strategy: **Allium (mainly autochory)**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Trophic mode

Parasitism and mycoheterotrophy: **autotrophic**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in the Czech Republic: **archaeophyte**
Invasion status: **casual**
Geographic origin: **anecophyte**
Period of introduction: **Eneolithic (4200–2300 BCE)**
Introduction pathway: **intentional – crops**

Habitat and sociology

Affinity to the forest environment

Affinity to the forest environment in Thermophyticum: **0 – taxon that does not spontaneously occur in Czech forests**
Affinity to the forest environment in Mesophyticum and Oreophyticum: **0 – taxon that does not spontaneously occur in Czech forests**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz