


Centaurea jacea × *C. nigrescens* × *C. oxylepis*

Distribution



Map info

● revised records

○ unrevised records

On the map are not visualized records without the coordinates and records marked as incorrect or doubtful.

Habitus and growth type

Life form: **hemicryptophyte**

Leaf

Leaf presence and metamorphosis: **leaves present, not modified**

Stipules: **absent**

Petiole: **both present and absent**

Flower

Flower colour: **pink-violet**

Perianth type: **calyx reduced, corolla present**

Calyx fusion: **pappus**

Inflorescence type: **corymbothyrsum ex anthodiis compositus**

Trophic mode

Parasitism and mycoheterotrophy: **autotrophic**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in the Czech Republic: **neophyte**

Invasion status: **casual**

Geographic origin: **hybrid origin**

Period of introduction: **Early Modern Period (1500-1800)**

Introduction pathway: **unintentional - nature**

Habitat and sociology

Affinity to the forest environment

Affinity to the forest environment in Thermophyticum: **0 - taxon that does not spontaneously occur in Czech forests**

Affinity to the forest environment in Mesophyticum and Oreophyticum: **0 - taxon that does not spontaneously occur in Czech forests**