


Festuca rupicola

Distribution



Map info

● revised records

○ unrevised records

On the map are not visualized records without the coordinates and records marked as incorrect or doubtful.



Habitus and growth type

Height [m]: **0.2-0.7**

Growth form: **clonal herb**

Life form: **hemicryptophyte**

Life strategy: **CS - competitor/stress-tolerator**

Life strategy (Pierce method based on leaf traits): **S/SR**

Life strategy (Pierce method, C-score): **7.2 %**

Life strategy (Pierce method, S-score): **78.5 %**

Life strategy (Pierce method, R-score): **14.4 %**

Leaf

Leaf presence and metamorphosis: **leaves present, not modified**

Leaf arrangement (phyllotaxis): **alternate**

Leaf shape: **simple - entire**

Stipules: **absent**

Petiole: **absent**

Leaf life span: **summer green**

Leaf anatomy: **scleromorphic**

Flower

Flowering period [month]: **May-June**



Flowering phase: **6 Cornus sanguinea-Melica uniflora (start of early summer)**
 Flower colour: **green**
 Perianth type: **reduced**
 Perianth fusion: **reduced**
 Inflorescence type: **panicula e spiculis composita**
 Dicliny: **synoecious**
 Pollination syndrome: **wind-pollination**

Fruit, seed and dispersal

Fruit type: **dry fruit - caryopsis**
 Fruit colour: **brown**
 Reproduction type: **mostly by seed/spores, rarely vegetatively**
 Dispersal unit (diaspore): **fruit, infrutescence or its part**
 Dispersal strategy: **Allium (mainly autochory)**
 Myrmecochory: **non-myrmecochorous (b)**

Belowground organs and clonality

Storage organ: **tuft**
 Type of clonal growth organ: **epigeogenous rhizome**
 Freely dispersible organs of clonal growth: **absent**
 Shoot life span (cyclicality): **dicyclic or polycyclic shoots prevailing**
 Branching type of stem-derived organs of clonal growth: **sympodial**
 Primary root: **absent**
 Persistence of the clonal growth organ [year]: **4**
 Number of clonal offspring: **3.5**
 Lateral spreading distance by clonal growth [m]: **0.01**
 Clonal index: **4**

Bud bank

Number of buds per shoot at the soil surface (root buds excluded): **5**
 Number of buds per shoot at a depth of 0–10 cm (root buds excluded): **15**
 Number of buds per shoot at a depth greater than 10 cm (root buds excluded): **0**
 Size of the belowground bud bank (root buds excluded): **20**
 Depth of the belowground bud bank (root buds excluded) [cm]: **4**
 Number of buds per shoot at the soil surface (root buds included): **5**
 Number of buds per shoot at a depth of 0–10 cm (root buds included): **15**
 Number of buds per shoot at a depth greater than 10 cm (root buds included): **0**
 Size of the belowground bud bank (root buds included): **20**
 Depth of the belowground bud bank (root buds included) [cm]: **4**

Trophic mode

Parasitism and mycoheterotrophy: **autotrophic**
 Carnivory: **non-carnivorous**
 Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**



Karyology

Chromosome number (2n): **42**

Ploidy level (x): **6**

2C genome size [Mbp]: **12331.29**

1Cx monoploid genome size [Mbp]: **2055.21**

Genomic GC content: **47.1 %**

Taxon origin

Origin in the Czech Republic: **native**

Ecological indicator values

Ellenberg-type indicator values

Light indicator value: **8 - light plant, only exceptionally occurring at less than 40% of diffuse radiation incident in an open area**

Temperature indicator value: **6 - transition between values 5 and 7**

Moisture indicator value: **3 - missing on damp soil**

Reaction indicator value: **8x - transition between values 7 and 9, occurring mostly in calcium-rich conditions (generalist)**

Nutrient indicator value: **3 - occurring at nutrient-poor sites more frequently than at average sites and exceptionally at rich sites**

Salinity indicator value: **0 - not salt tolerant, glycophyte**

Indicator values for disturbance

Whole-community disturbance frequency indicator value: **-0.71**

Herb layer disturbance frequency indicator value: **-0.35**

Whole-community disturbance severity indicator value: **0.25**

Herb layer disturbance severity indicator value: **0.28**

Whole-community structure based disturbance indicator value: **0.49**

Herb layer structure-based disturbance indicator value: **0.57**

Habitat and sociology

Occurrence in habitats

1 Vegetation of cliffs, screes and walls

1A Calcareous cliffs: **1 - rare occurrence**

1B Siliceous cliffs and block fields: **1 - rare occurrence**

1D Mobile calcareous screes: **1 - rare occurrence**

6 Meadows and mesic pastures

6A Mesic Arrhenatherum meadows: **2 - optimum**

6C Pastures and park grasslands: **1 - rare occurrence**

6D Alluvial meadows of lowland rivers: **1 - rare occurrence**



6F Intermittently wet Molinia meadows: **1 - rare occurrence**

7 Acidophilous grasslands

7B Submontane Nardus grasslands: **1 - rare occurrence**

8 Dry grasslands

8A Hercynian dry grasslands on rock outcrops: **1 - rare occurrence**



- 8B Submediterranean dry grasslands on rock outcrops: **2 - optimum**
- 8C Narrow-leaved sub-continental steppes: **3 - dominant**
- 8D Broad-leaved dry grasslands: **2 - optimum**
- 8E Acidophilous dry grasslands: **3 - dominant**
- 8F Thermophilous forest fringe vegetation: **2 - optimum**
- 9 Sand grasslands and rock-outcrop vegetation
- 9B Open vegetation of acidic sands: **1 - rare occurrence**
- 9C Festuca grasslands on acidic sands: **2 - optimum**
- 9D Pannonian sand steppes: **1 - rare occurrence**
- 9E Acidophilous vegetation of spring therophytes and succulents: **1 - rare occurrence**
- 9F Basiphilous vegetation of spring therophytes and succulents: **2 - optimum**
- 10 Saline vegetation
- 10I Inland saline meadows: **1 - rare occurrence**
- 11 Heathlands and scrub
- 11A Dry lowland to subalpine heathlands: **1 - rare occurrence**
- 11L Tall mesic and xeric shrub: **1 - rare occurrence**
- 11N Low xeric scrub: **2 - optimum**
- 12 Forests
- 12C Oak-hornbeam forests: **1 - rare occurrence**
- 12D Ravine forests: **1 - rare occurrence**
- 12F Limestone beech forests: **1 - rare occurrence**
- 12H Peri-Alpidic basiphilous thermophilous oak forests: **2 - optimum**
- 12I Sub-continental thermophilous oak forests: **1 - rare occurrence**
- 12J Acidophilous thermophilous oak forests: **1 - rare occurrence**
- 12K Acidophilous oak forests: **1 - rare occurrence**
- 12L Boreo-continental pine forests: **1 - rare occurrence**
- 12O Peri-Alpidic pine forests: **2 - optimum**
- 12T Robinia pseudacacia plantations: **1 - rare occurrence**
- 12U Plantations of broad-leaved non-native trees: **1 - rare occurrence**
- 12W Pine and larch plantations: **1 - rare occurrence**
- 13 Anthropogenic vegetation
- 13C Annual vegetation of trampled habitats: **1 - rare occurrence**
- 13D Perennial thermophilous ruderal vegetation: **1 - rare occurrence**
- Affinity to the forest environment
- Affinity to the forest environment in Thermophyticum: **0 - taxon that does not spontaneously occur in Czech forests**
- Affinity to the forest environment in Mesophyticum and Oreophyticum: **0 - taxon that does not spontaneously occur in Czech forests**
- Diagnostic taxon
- Diagnostic taxon of classes: [TH Festuco-Brometea](#)
- Diagnostic taxon of alliances: [THE Cirsio-Brachypodium pinnati](#), [THF Bromion erecti](#)
- Diagnostic taxon of associations: [THD03 Festuco rupicolae-Caricetum humilis](#), [THD06 Astragalo exscapi-Crambetum tatariae](#), [THE01 Scabioso ochroleucae-Brachypodietum pinnati](#), [THE03 Polygalo majoris-Brachypodietum pinnati](#), [THF02 Brachypodio pinnati-Molinietum arundinaceae](#), [XCA02 Salvia nemorosae-Marrubietum peregrini](#)

Constant taxon

Constant taxon of classes: [TH Festuco-Brometea](#)

Constant taxon of alliances: [THD Festucion valesiacae](#), [THE Cirsio-Brachypodion pinnati](#), [THF Bromion erecti](#)

Constant taxon of associations: [LCA02 Lithospermo purpureocaerulei-Quercetum pubescentis](#), [TFC01 Sileno otitae-Festucetum brevipilae](#), [THC04 Asplenio cuneifolii-Seslerietum caeruleae](#), [THD03 Festuco rupicolae-Caricetum humilis](#), [THD04 Koelerio macranthae-Stipetum joannis](#), [THD05 Stipetum tirsae](#), [THD06 Astragalo excapici-Crambetum tatariae](#), [THE01 Scabioso ochroleucae-Brachypodietum pinnati](#), [THE03 Polygalo majoris-Brachypodietum pinnati](#), [THE04 Plantagini maritimae-Caricetum flaccae](#), [THF01 Carlino acaulis-Brometum erecti](#), [THF02 Brachypodio pinnati-Molinietum arundinaceae](#), [THG01 Potentillo heptaphyllae-Festucetum rupicolae](#), [THH03 Geranio sanguinei-Peucedanetum cervariae](#), [XCA02 Salvia nemorosae-Marrubietum peregrini](#)

Dominant taxon

Dominant taxon of associations: [KBB04 Pruno spinosae-Ligustretum vulgaris](#), [TDA04 Potentillo albae-Festucetum rubrae](#), [THD01 Festuco valesiacae-Stipetum capillatae](#), [THD03 Festuco rupicolae-Caricetum humilis](#), [THE01 Scabioso ochroleucae-Brachypodietum pinnati](#), [THF01 Carlino acaulis-Brometum erecti](#), [THG01 Potentillo heptaphyllae-Festucetum rupicolae](#), [XCA02 Salvia nemorosae-Marrubietum peregrini](#)

Ecological specialization indices

Ecological specialization index for all vegetation types: **4.6**

Ecological specialization index for non-forest vegetation: **4.6**

Ecological specialization index for forest vegetation: **4.8**

Colonization ability

Index of colonization success (ICS): **8**

Index of colonization potential (ICP): **5**

Optimum successional age [years]: **32**

Distribution and frequency

Floristic zone: **northern temperate, southern temperate, submeridional, meridional**

Floristic region: **Europe, Western Asia**

Continental degree: **8**

Distribution range extension along the continentality gradient: **4**

Elevational belt in the Czech Republic: **lowlands, colline belt, submontane belt**

Occurrence frequency in the basic grid mapping cells and quadrants of the basic grid mapping cells: 464

taxon.data.freq_in_quad: 1228

Commonness in vegetation plots from the Czech Republic

Occurrence frequency in vegetation plots: **3.6 %**

Occurrence frequency in vegetation plots with a cover above 5%: **38 %**

Occurrence frequency in vegetation plots with a cover above 25%: **13.2 %**

Occurrence frequency in vegetation plots with a cover above 50%: **3.5 %**

Mean percentage cover in vegetation plots: **10.6 %**

Maximum percentage cover in vegetation plots: **63 %**

Number of habitats with taxon occurrence in the Czech Republic

Number of narrow habitats in which the taxon occurs: **37**

Number of narrow habitats in which the taxon has its optimum: **11**

Number of broad habitats in which the taxon occurs: **9**

Number of broad habitats in which the taxon has its optimum: **5**

Threats and protection

Red List 2017 (national categories): **taxon is not on the Red List**

Red List 2017 (IUCN categories): **LC(NA) - least concern (taxon is not on the Red List)**

Legal protection: **not protected by law**