
Cochlearia danica

Distribution



Habitus and growth type

Height [m]: **0.1–0.2**
Life form: **therophyte (hemicryptophyte)**
Life strategy (Pierce method based on leaf traits): **R**
Life strategy (Pierce method, C-score): **9.6 %**
Life strategy (Pierce method, S-score): **0 %**
Life strategy (Pierce method, R-score): **90.4 %**

Flower

Flowering period [month]: **May–June**
Flower colour: **white**
Inflorescence type: **racemus**

Trophic mode

Parasitism and mycoheterotrophy: **autotrophic**
Carnivory: **non-carnivorous**

Taxon origin

Origin in the Czech Republic: **neophyte**
Invasion status: **invasive**
Geographic origin: **Europe**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Year of the first record in the wild: **2016**
Period of introduction: **Present (after 2000)**
Introduction pathway: **unintentional – traffic**

Threats and protection

Red List 2017 (national categories): **taxon is not on the Red List**
Red List 2017 (IUCN categories): **LC(NA) – least concern (taxon is not on the Red List)**
Legal protection: **not protected by law**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz