
Crataegus laevigata agg.

Distribution





Map info

revised records

unrevised records

On the map are not visualized records without the coordinates and records marked as incorrect or doubtful.





Habitus and growth type

Height [m]: **1–10**
Growth form: **shrub (tree)**
Life form: **macrophanerophyte, nanophanerophyte**
Life strategy: **C – competitor**

Leaf

Leaf presence and metamorphosis: **leaves present, not modified**
Leaf arrangement (phyllotaxis): **alternate**
Leaf shape: **simple – pinnately divided**
Stipules: **present**
Petiole: **present**
Leaf life span: **summer green**
Leaf deciduousness in woody plants: **winter deciduous**
Leaf anatomy: **mesomorphic**
Functional leaf type in woody plants: **broad deciduous or semi-deciduous**

Flower

Flowering period [month]: **May–June**
Flowering phase: **5 Sorbus aucuparia-Galium odoratum (end of mid-spring)**
Flower colour: **white**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Flower symmetry: **actinomorphic**
Perianth type: **calyx and corolla**
Perianth fusion: **free**
Calyx fusion: **hypanthium**
Inflorescence type: **corymbus**
Dicliny: **synoecious**
Generative reproduction type: **facultative allogamy**
Pollination syndrome: **insect-pollination, selfing**



Fruit, seed and dispersal

Fruit type: **fleshy fruit – pome**
Fruit colour: **red**
Reproduction type: **only by seed/spores**
Dispersal unit (diaspore): **seed, fruit, infrutescence or its part**
Dispersal strategy: **Cornus (mainly autochory and endozoochory)**
Myrmecochory: **non-myrmecochorous (b)**



Belowground organs and clonality

Shoot metamorphosis: **shoot thorn**

Bud bank

Number of buds per shoot at the soil surface (root buds excluded): **0**
Number of buds per shoot at a depth of 0–10 cm (root buds excluded): **5**
Number of buds per shoot at a depth greater than 10 cm (root buds excluded): **5**
Size of the belowground bud bank (root buds excluded): **10**
Depth of the belowground bud bank (root buds exluded) [cm]: **10**
Number of buds per shoot at the soil surface (root buds included): **0**
Number of buds per shoot at a depth of 0–10 cm (root buds included): **5**
Number of buds per shoot at a depth greater than 10 cm (root buds included): **5**
Size of the belowground bud bank (root buds included): **10**
Depth of the belowground bud bank (root buds included) [cm]: **10**



Trophic mode

Parasitism and mycoheterotrophy: **autotrophic**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**



Karyology

Chromosome number (2n): **34, 68**
Ploidy level (x): **2, 4**
2C genome size [Mbp]: **1961.95**
1Cx monoploid genome size [Mbp]: **654**



Taxon origin

Origin in the Czech Republic: **native**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Ecological indicator values

Ellenberg-type indicator values

Light indicator value: **5x – semi-shade plant, only exceptionally occurring in full light, but usually at more than 10% of the diffuse radiation incident in an open area (generalist)**

Temperature indicator value: **6 – transition between values 5 and 7**

Moisture indicator value: **5 – indicator of fresh soils, focus on soils of average moisture, missing on wet and on soils that frequently dry out**

Reaction indicator value: **7 – indicator of slightly acidic to slightly basic conditions, never occurring in very acidic conditions**

Nutrient indicator value: **5 – occurring at moderately nutrient-rich sites, and less frequently at poor and rich sites**

Salinity indicator value: **0 – not salt tolerant, glycophyte**

Habitat and sociology

Occurrence in habitats

8 Dry grasslands

8A Hercynian dry grasslands on rock outcrops: **1 – rare occurrence**

8B Submediterranean dry grasslands on rock outcrops: **1 – rare occurrence**

8C Narrow-leaved sub-continental steppes: **1 – rare occurrence**

8D Broad-leaved dry grasslands: **1 – rare occurrence**

8E Acidophilous dry grasslands: **1 – rare occurrence**

8F Thermophilous forest fringe vegetation: **1 – rare occurrence**

11 Heathlands and scrub

11I Willow carrs: **1 – rare occurrence**

11J Willow galleries of loamy and sandy river banks: **1 – rare occurrence**

11L Tall mesic and xeric shrub: **2 – optimum**

11N Low xeric scrub: **1 – rare occurrence**

11R Scrub and pioneer woodland of forests clearings: **1 – rare occurrence**

12 Forests

12A Alder carrs: **1 – rare occurrence**

12B Alluvial forests: **2 – optimum**

12C Oak-hornbeam forests: **2 – optimum**

12D Ravine forests: **1 – rare occurrence**

12E Herb-rich beech forests: **1 – rare occurrence**

12F Limestone beech forests: **1 – rare occurrence**

12H Peri-Alpidic basiphilous thermophilous oak forests: **2 – optimum**

12I Sub-continental thermophilous oak forests: **2 – optimum**

12J Acidophilous thermophilous oak forests: **1 – rare occurrence**

12K Acidophilous oak forests: **1 – rare occurrence**

12O Peri-Alpidic pine forests: **1 – rare occurrence**

12T Robinia pseudacacia plantations: **1 – rare occurrence**

12U Plantations of broad-leaved non-native trees: **1 – rare occurrence**

12W Pine and larch plantations: **1 – rare occurrence**

Affinity to the forest environment

Affinity to the forest environment in Thermophyticum: **2.1 – taxon occurring both in**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

the forest and open vegetation
Affinity to the forest environment in Mesophyticum and Oreophyticum: **2.1 – taxon occurring both in the forest and open vegetation**

Distribution and frequency

Floristic zone: **northern temperate, southern temperate, submeridional**
Floristic region: **Europe**
Continentality degree: **5**
Distribution range extension along the continentality gradient: **3**
Elevational belt in the Czech Republic: **lowlands, colline belt, submontane belt, montane belt**
Expansive taxon in the region: **Bohemian Thermophyticum, Bohemian Moravian Mesophyticum, Pannonian Thermophyticum, Carpathian Mesophyticum**
Occurrence frequency in the basic grid mapping cells and quadrants of the basic grid mapping cells: 600
taxon.data.freq_in_quad: 1637

Commonness in vegetation plots from the Czech Republic

Maximum percentage cover in vegetation plots: **46 %**

Number of habitats with taxon occurrence in the Czech Republic

Number of narrow habitats in which the taxon occurs: **25**
Number of narrow habitats in which the taxon has its optimum: **5**
Number of broad habitats in which the taxon occurs: **3**
Number of broad habitats in which the taxon has its optimum: **2**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz