
Acer tataricum

Distribution





Map info

revised records

unrevised records

On the map are not visualized records without the coordinates and records marked as incorrect or doubtful.









Habitus and growth type

Height [m]: **2–8**
Growth form: **shrub (tree)**
Life form: **nanophanerophyte (macrophanerophyte)**
Life strategy (Pierce method based on leaf traits): **S/CSR**
Life strategy (Pierce method, C-score): **25.1 %**
Life strategy (Pierce method, S-score): **54.8 %**
Life strategy (Pierce method, R-score): **20.1 %**

Leaf

Leaf presence and metamorphosis: **leaves present, not modified**
Leaf arrangement (phyllotaxis): **opposite**
Leaf shape: **simple – palmately divided**
Stipules: **absent**
Petiole: **present**
Leaf deciduousness in woody plants: **winter deciduous**
Functional leaf type in woody plants: **broad deciduous or semi-deciduous**

Flower

Flowering period [month]: **May–June**
Flower colour: **green-white**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz



Flower symmetry: **actinomorphic**
Perianth type: **calyx and corolla**
Perianth fusion: **free**
Calyx fusion: **aposepalous**
Inflorescence type: **panicula**

Fruit, seed and dispersal

Fruit type: **dry fruit – double samara**
Fruit colour: **red, brown**
Dispersal unit (diaspore): **fruit, infrutescence or its part**
Dispersal strategy: **Epilobium (mainly anemochory and autochory)**
Myrmecochory: **non-myrmecochorous (b)**



Trophic mode

Parasitism and mycoheterotrophy: **autotrophic**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin



Origin in the Czech Republic: **neophyte**
Invasion status: **naturalized**
Geographic origin: **Europe**
Year of the first record in the wild: **2004**
Period of introduction: **Present (after 2000)**
Introduction pathway: **intentional – ornamental, intentional – landscaping**

Habitat and sociology

Affinity to the forest environment

Affinity to the forest environment in Thermophyticum: **0 – taxon that does not spontaneously occur in Czech forests**
Affinity to the forest environment in Mesophyticum and Oreophyticum: **0 – taxon that does not spontaneously occur in Czech forests**

Threats and protection

Red List 2017 (national categories): **taxon is not on the Red List**
Red List 2017 (IUCN categories): **LC(NA) – least concern (taxon is not on the Red List)**
Legal protection: **not protected by law**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz