
Ajuga genevensis

Distribution





Map info

revised records

unrevised records

On the map are not visualized records without the coordinates and records marked as incorrect or doubtful.







Habitus and growth type

Height [m]: **0.1–0.3**
Growth form: **clonal herb**
Life form: **hemicryptophyte (geophyte)**
Life strategy: **CSR – competitor/stress-tolerator/ruderal**
Life strategy (Pierce method based on leaf traits): **R/CR**
Life strategy (Pierce method, C-score): **24.9 %**
Life strategy (Pierce method, S-score): **0 %**
Life strategy (Pierce method, R-score): **75.1 %**

Leaf

Leaf presence and metamorphosis: **leaves present, not modified**
Leaf arrangement (phyllotaxis): **opposite, rosulate**
Leaf shape: **simple – entire**
Stipules: **absent**
Petiole: **both present and absent**
Leaf life span: **summer green**
Leaf anatomy: **mesomorphic**

Flower

Flowering period [month]: **April–July**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Flowering phase: **5 Sorbus aucuparia-Galium odoratum (end of mid-spring)**
Flower colour: **white, pink, blue-violet**
Flower symmetry: **zygomorphic**
Perianth type: **calyx and corolla**
Perianth fusion: **fused**
Shape of the sympetalous corolla or syntepalous perianth: **bilabiate**
Calyx fusion: **synsepalous**
Inflorescence type: **pseudospica e verticillastris composita**
Dicliny: **synoecious**
Generative reproduction type: **facultative allogamy**
Pollination syndrome: **insect-pollination, selfing**



Fruit, seed and dispersal

Fruit type: **dry fruit – cluster of four one-seeded nutlets**
Fruit colour: **brown**
Reproduction type: **mostly by seed/spores, rarely vegetatively**
Dispersal unit (diaspore): **seed**
Dispersal strategy: **Allium (mainly autochory)**
Myrmecochory: **myrmecochorous**





Belowground organs and clonality

Root metamorphosis: **root shoot**
Type of clonal growth organ: **root with adventitious buds**
Freely dispersible organs of clonal growth: **absent**
Shoot life span (cyclicity): **dicyclic or polycyclic shoots prevailing**
Branching type of stem-derived organs of clonal growth: **sympodial**
Primary root: **absent**
Persistence of the clonal growth organ [year]: **2.7**
Number of clonal offspring: **4**
Lateral spreading distance by clonal growth [m]: **0.1**
Clonal index: **5**
Position of root buds: **lateral roots**
Role of root buds in life-history of a plant: **additive**






Bud bank

Number of buds per shoot at the soil surface (root buds excluded): **5**
Number of buds per shoot at a depth of 0–10 cm (root buds excluded): **12**
Number of buds per shoot at a depth greater than 10 cm (root buds excluded): **0**
Size of the belowground bud bank (root buds excluded): **17**
Depth of the belowground bud bank (root buds exluded) [cm]: **4**
Number of buds per shoot at the soil surface (root buds included): **5**
Number of buds per shoot at a depth of 0–10 cm (root buds included): **20**
Number of buds per shoot at a depth greater than 10 cm (root buds included): **15**
Size of the belowground bud bank (root buds included): **40**
Depth of the belowground bud bank (root buds included) [cm]: **8**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz





Trophic mode

Parasitism and mycoheterotrophy: **autotrophic**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Karyology

Chromosome number (2n): **32**
Ploidy level (x): **4**
2C genome size [Mbp]: **2228.81**
1Cx monoploid genome size [Mbp]: **557.2**
Genomic GC content: **40 %**

Taxon origin

Origin in the Czech Republic: **native**

Ecological indicator values

Ellenberg-type indicator values

Light indicator value: **7 – half-light plant, mostly occurring at full light, but also in the shade up to about 30% of diffuse radiation incident in an open area**
Temperature indicator value: **6x – transition between values 5 and 7 (generalist)**
Moisture indicator value: **4 – transition between values 3 and 5**
Reaction indicator value: **7 – indicator of slightly acidic to slightly basic conditions, never occurring in very acidic conditions**
Nutrient indicator value: **3 – occurring at nutrient-poor sites more frequently than at average sites and exceptionally at rich sites**
Salinity indicator value: **0 – not salt tolerant, glycophyte**

Indicator values for disturbance

Whole-community disturbance frequency indicator value: **-1.59**
Herb layer disturbance frequency indicator value: **-0.57**
Whole-community disturbance severity indicator value: **0.25**
Herb layer disturbance severity indicator value: **0.2**
Whole-community structure based disturbance indicator value: **0.27**
Herb layer structure-based disturbance indicator value: **0.35**

Habitat and sociology

Occurrence in habitats

1 Vegetation of cliffs, screes and walls

1D Mobile calcareous screes: **1 – rare occurrence**

6 Meadows and mesic pastures

6A Mesic Arrhenatherum meadows: **1 – rare occurrence**
6C Pastures and park grasslands: **1 – rare occurrence**
6F Intermittently wet Molinia meadows: **1 – rare occurrence**

7 Acidophilous grasslands

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

7B Submontane Nardus grasslands: **1 – rare occurrence**

8 Dry grasslands

8A Hercynian dry grasslands on rock outcrops: **2 – optimum**
8B Submediterranean dry grasslands on rock outcrops: **1 – rare occurrence**
8C Narrow-leaved sub-continental steppes: **1 – rare occurrence**
8D Broad-leaved dry grasslands: **1 – rare occurrence**
8E Acidophilous dry grasslands: **2 – optimum**
8F Thermophilous forest fringe vegetation: **2 – optimum**

9 Sand grasslands and rock-outcrop vegetation

9E Acidophilous vegetation of spring therophytes and succulents: **1 – rare occurrence**
9F Basiphilous vegetation of spring therophytes and succulents: **1 – rare occurrence**

11 Heathlands and scrub

11A Dry lowland to subalpine heathlands: **1 – rare occurrence**
11L Tall mesic and xeric shrub: **1 – rare occurrence**
11N Low xeric scrub: **1 – rare occurrence**
11R Scrub and pioneer woodland of forests clearings: **1 – rare occurrence**

12 Forests

12C Oak-hornbeam forests: **1 – rare occurrence**
12D Ravine forests: **1 – rare occurrence**
12F Limestone beech forests: **2 – optimum**
12G Acidophilous beech forests: **1 – rare occurrence**
12H Peri-Alpidic basiphilous thermophilous oak forests: **2 – optimum**
12I Sub-continental thermophilous oak forests: **2 – optimum**
12J Acidophilous thermophilous oak forests: **2 – optimum**
12K Acidophilous oak forests: **1 – rare occurrence**
12L Boreo-continental pine forests: **1 – rare occurrence**
12O Peri-Alpidic pine forests: **1 – rare occurrence**
12V Spruce plantations: **1 – rare occurrence**
12W Pine and larch plantations: **1 – rare occurrence**

13 Anthropogenic vegetation

13F Herbaceous vegetation of forests clearings and Rubus scrub: **2 – optimum**

Affinity to the forest environment

Affinity to the forest environment in Thermophyticum: **2.1 – taxon occurring both in the forest and open vegetation**
Affinity to the forest environment in Mesophyticum and Oreophyticum: **2.2 – taxon occurring partly in the forest, but mainly in open vegetation**

Diagnostic taxon

Diagnostic taxon of alliances: **LCA *Quercion pubescenti-petraeae*, LCB *Aceri tatarici-Quercion***
Diagnostic taxon of associations: **LCA02 *Lithospermo purpurocaerulei-Quercetum pubescentis***

Constant taxon

Constant taxon of associations: **LCA02 *Lithospermo purpurocaerulei-Quercetum pubescentis***

Ecological specialization indices

Ecological specialization index for all vegetation types: **3.9**
Ecological specialization index for non-forest vegetation: **4.2**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Ecological specialization index for forest vegetation: **4.5**

Colonization ability

Index of colonization success (ICS): **5**
Index of colonization potential (ICP): **5**
Optimum successional age [years]: **13**

Distribution and frequency

Floristic zone: **northern temperate, southern temperate, submeridional, meridional**
Floristic region: **Europe**
Continentality degree: **5**
Distribution range extension along the continentality gradient: **4**
Elevational belt in the Czech Republic: **lowlands, colline belt (submontane belt)**
Occurrence frequency in the basic grid mapping cells and quadrants of the basic grid mapping cells: 534
taxon.data.freq_in_quad: 1374

Commonness in vegetation plots from the Czech Republic

Occurrence frequency in vegetation plots: **0.6 %**
Occurrence frequency in vegetation plots with a cover above 5%: **1.6 %**
Occurrence frequency in vegetation plots with a cover above 25%: **0 %**
Occurrence frequency in vegetation plots with a cover above 50%: **0 %**
Mean percentage cover in vegetation plots: **2.1 %**
Maximum percentage cover in vegetation plots: **13 %**

Number of habitats with taxon occurrence in the Czech Republic

Number of narrow habitats in which the taxon occurs: **30**
Number of narrow habitats in which the taxon has its optimum: **8**
Number of broad habitats in which the taxon occurs: **8**
Number of broad habitats in which the taxon has its optimum: **3**

Threats and protection

Red List 2017 (national categories): **taxon is not on the Red List**
Red List 2017 (IUCN categories): **LC(NA) – least concern (taxon is not on the Red List)**
Legal protection: **not protected by law**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz