
Amelanchier lamarckii

Distribution





Map info

revised records

unrevised records

On the map are not visualized records without the coordinates and records marked as incorrect or doubtful.





Habitus and growth type

Height [m]: **2–10**
Growth form: **shrub (tree)**
Life form: **nanophanerophyte (macrophanerophyte)**
Life strategy (Pierce method based on leaf traits): **S/CS**
Life strategy (Pierce method, C-score): **26.4 %**
Life strategy (Pierce method, S-score): **67 %**
Life strategy (Pierce method, R-score): **6.6 %**


Leaf

Leaf presence and metamorphosis: **leaves present, not modified**
Leaf arrangement (phyllotaxis): **alternate**
Leaf shape: **simple – entire**
Stipules: **present**
Petiole: **present**
Functional leaf type in woody plants: **broad deciduous or semi-deciduous**

Flower

Flowering period [month]: **April–May**
Flower colour: **white**
Flower symmetry: **actinomorphic**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz







Perianth type: **calyx and corolla**
Perianth fusion: **free**
Calyx fusion: **hypanthium**
Inflorescence type: **racemus**
Generative reproduction type: **apomixis**

Fruit, seed and dispersal

Fruit type: **fleshy fruit – pome**
Dispersal unit (diaspore): **seed, fruit, infrutescence or its part**
Dispersal strategy: **Cornus (mainly autochory and endozoochory)**
Myrmecochory: **non-myrmecochorous (b)**

Trophic mode

Parasitism and mycoheterotrophy: **autotrophic**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in the Czech Republic: **neophyte**
Invasion status: **naturalized**
Geographic origin: **North America**
Year of the first record in the wild: **1867**
Period of introduction: **Late Modern Period (1800–1950)**
Introduction pathway: **intentional – ornamental, intentional – nature**

Ecological indicator values

Ellenberg-type indicator values

Light indicator value: **6 – transition between values 5 and 7; rarely at less than 20% of diffuse radiation incident in an open area**
Temperature indicator value: **6 – transition between values 5 and 7**
Moisture indicator value: **5 – indicator of fresh soils, focus on soils of average moisture, missing on wet and on soils that frequently dry out**
Reaction indicator value: **3 – acidity indicator, occurring mainly in acidic conditions, exceptionally in neutral conditions**
Nutrient indicator value: **3 – occurring at nutrient-poor sites more frequently than at average sites and exceptionally at rich sites**
Salinity indicator value: **0 – not salt tolerant, glycophyte**

Habitat and sociology

Affinity to the forest environment

Affinity to the forest environment in Thermophyticum: **0 – taxon that does not spontaneously occur in Czech forests**
Affinity to the forest environment in Mesophyticum and Oreophyticum: **0 – taxon that does not spontaneously occur in Czech forests**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Distribution and frequency

Continentality degree: **5**
Distribution range extension along the continentality gradient: **4**
Occurrence frequency in the basic grid mapping cells and quadrants of the basic grid mapping cells: 18
taxon.data.freq_in_quad: 21

Threats and protection

Red List 2017 (national categories): **taxon is not on the Red List**
Red List 2017 (IUCN categories): **LC(NA) – least concern (taxon is not on the Red List)**
Legal protection: **not protected by law**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz