
Cardamine amara

Distribution





Map info

revised records

unrevised records

On the map are not visualized records without the coordinates and records marked as incorrect or doubtful.







Habitus and growth type

Height [m]: **0.1–0.6**
Growth form: **clonal herb**
Life form: **hemicryptophyte**
Life strategy: **CSR – competitor/stress-tolerator/ruderal**
Life strategy (Pierce method based on leaf traits): **CR**
Life strategy (Pierce method, C-score): **41.6 %**
Life strategy (Pierce method, S-score): **0 %**
Life strategy (Pierce method, R-score): **58.4 %**

Leaf

Leaf presence and metamorphosis: **leaves present, not modified**
Leaf arrangement (phyllotaxis): **alternate**
Leaf shape: **compound – imparipinnate**
Stipules: **absent**
Petiole: **present**
Leaf life span: **evergreen**
Leaf anatomy: **hygromorphic, helomorphic**

Flower

Flowering period [month]: **April–July**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Flowering phase: **5 Sorbus aucuparia-Galium odoratum (end of mid-spring)**
Flower colour: **white**
Flower symmetry: **actinomorphic**
Perianth type: **calyx and corolla**
Perianth fusion: **free**
Calyx fusion: **aposepalous**
Inflorescence type: **racemus**
Dicliny: **synoecious, gynomonoecious**
Generative reproduction type: **facultative allogamy**
Pollination syndrome: **insect-pollination, selfing**




Fruit, seed and dispersal



Fruit type: **dry fruit – siliqua**
Fruit colour: **brown**
Reproduction type: **by seed/spores and vegetatively**
Dispersal unit (diaspore): **seed, shoot fragment**
Dispersal strategy: **Sparganium (mainly autochory and hydrochory)**
Myrmecochory: **non-myrmecochorous (b)**

Belowground organs and clonality



Shoot metamorphosis: **stolon, rhizome**
Storage organ: **stolon, rhizome**
Type of clonal growth organ: **stolon**
Freely dispersible organs of clonal growth: **absent**
Shoot life span (cyclicity): **dicyclic or polycyclic shoots prevailing**
Branching type of stem-derived organs of clonal growth: **sympodial**
Primary root: **absent**
Persistence of the clonal growth organ [year]: **1.5**
Number of clonal offspring: **1**
Lateral spreading distance by clonal growth [m]: **0.13**
Clonal index: **4**

Bud bank



Number of buds per shoot at the soil surface (root buds excluded): **15**
Number of buds per shoot at a depth of 0–10 cm (root buds excluded): **0**
Number of buds per shoot at a depth greater than 10 cm (root buds excluded): **0**
Size of the belowground bud bank (root buds excluded): **15**
Depth of the belowground bud bank (root buds exluded) [cm]: **1**
Number of buds per shoot at the soil surface (root buds included): **15**
Number of buds per shoot at a depth of 0–10 cm (root buds included): **0**
Number of buds per shoot at a depth greater than 10 cm (root buds included): **0**
Size of the belowground bud bank (root buds included): **15**
Depth of the belowground bud bank (root buds included) [cm]: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotrophic**
Carnivory: **non-carnivorous**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**



Karyology

Chromosome number (2n): **16, 32**
Ploidy level (x): **2, 4**
2C genome size [Mbp]: **717.83**
1Cx monoploid genome size [Mbp]: **214.59**

Taxon origin

Origin in the Czech Republic: **native**



Ecological indicator values

Ellenberg-type indicator values

Light indicator value: **5 – semi-shade plant, only exceptionally occurring in full light, but usually at more than 10% of the diffuse radiation incident in an open area**
Temperature indicator value: **5 – moderate heat indicator, occurring from lowland to montane belt, mainly in submontane-temperate areas**
Moisture indicator value: **9 – wetness indicator, focus on often soaked, poorly aerated soils**
Reaction indicator value: **6 – transition between values 5 and 7**
Nutrient indicator value: **6 – transition between values 5 and 7**
Salinity indicator value: **0 – not salt tolerant, glycophyte**



Habitat and sociology

Occurrence in habitats

1 Vegetation of cliffs, screes and walls

1B Siliceous cliffs and block fields: **1 – rare occurrence**

2 Alpine and subalpine grasslands

2B Subalpine tall-forb and tall-grass vegetation: **1 – rare occurrence**

3 Aquatic vegetation

3B Macrophytic vegetation of water streams: **1 – rare occurrence**



4 Wetland and riverine herbaceous vegetation

4A Reed-beds of eutrophic still waters: **1 – rare occurrence**
4C Eutrophic vegetation of muddy substrata: **1 – rare occurrence**
4D Riverine reed vegetation: **1 – rare occurrence**
4E Reed vegetation of brooks: **2 – optimum**
4F Mesotrophic vegetation of muddy substrata: **1 – rare occurrence**
4G Tall-sedge beds: **1 – rare occurrence**
4H Vegetation of low annual hygrophilous herbs: **1 – rare occurrence**
4I Vegetation of nitrophilous annual hygrophilous herbs: **1 – rare occurrence**
4J River gravel banks: **1 – rare occurrence**
4K Petasites fringes of montane brooks: **2 – optimum**



5 Vegetation of springs and mires







5A Hard-water springs with tufa formation: **2 – optimum**
5B Lowland to montane soft-water springs: **3 – dominant**
5C Alpine and subalpine soft-water springs: **3 – dominant**
5D Calcareous fens: **1 – rare occurrence**
5E Acidic moss-rich fens and peatland meadows: **1 – rare occurrence**
5F Transitional mires: **1 – rare occurrence**

6 Meadows and mesic pastures

6E Wet Cirsium meadows: **2 – optimum**
6G Vegetation of wet disturbed soils: **1 – rare occurrence**

11 Heathlands and scrub

11H Subalpine deciduous scrub: **1 – rare occurrence**
11I Willow carrs: **1 – rare occurrence**
11J Willow galleries of loamy and sandy river banks: **2 – optimum**

12 Forests

12A Alder carrs: **2 – optimum**
12B Alluvial forests: **2 – optimum**
12E Herb-rich beech forests: **1 – rare occurrence**
12G Acidophilous beech forests: **1 – rare occurrence**
12R Acidophilous spruce forests: **1 – rare occurrence**
12V Spruce plantations: **1 – rare occurrence**

13 Anthropogenic vegetation

13E Perennial nitrophilous herbaceous vegetation of mesic sites: **1 – rare occurrence**

Diagnostic taxon

Diagnostic taxon of classes: **LA *Alnetea glutinosae***, **RA *Montio-Cardaminetea***
Diagnostic taxon of alliances: **KAB *Salicion elaeagno-daphnoidis***, **LAA *Alnion glutinosae***, **RAA *Caricion remotae***
Diagnostic taxon of associations: **KAB02 *Salicetum purpureae***, **LAA03 *Carici acutiformis-Alnetum glutinosae***, **LBA03 *Carici remotae-Fraxinetum excelsioris***, **LFC03 *Equiseto sylvatici-Piceetum abietis***, **RAA02 *Cardamino-Chrysosplenietum alternifolii***, **RAA03 *Pellio epiphyllae-Chrysosplenietum oppositifolii***

Constant taxon

Constant taxon of classes: **RA *Montio-Cardaminetea***
Constant taxon of alliances: **RAA *Caricion remotae***
Constant taxon of associations: **KAB02 *Salicetum purpureae***, **LAA03 *Carici acutiformis-Alnetum glutinosae***, **LBA03 *Carici remotae-Fraxinetum excelsioris***, **LFC03 *Equiseto sylvatici-Piceetum abietis***, **RAA02 *Cardamino-Chrysosplenietum alternifolii***, **RAA03 *Pellio epiphyllae-Chrysosplenietum oppositifolii***

Dominant taxon

Dominant taxon of associations: **LBA03 *Carici remotae-Fraxinetum excelsioris***, **RAA02 *Cardamino-Chrysosplenietum alternifolii***, **RAA03 *Pellio epiphyllae-Chrysosplenietum oppositifolii***

Colonization ability

Index of colonization success (ICS): **4**
Index of colonization potential (ICP): **1**
Optimum successional age [years]: **1**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Distribution and frequency

Floristic zone: **boreal, northern temperate, southern temperate, submeridional**
Floristic region: **Europe, Western Siberia**
Continentality degree: **5**
Distribution range extension along the continentality gradient: **4**
Elevational belt in the Czech Republic: **lowlands, colline belt, submontane belt, montane belt, subalpine belt**
Occurrence frequency in the basic grid mapping cells and quadrants of the basic grid mapping cells: 638
taxon.data.freq_in_quad: 2151

Commonness in vegetation plots from the Czech Republic

Maximum percentage cover in vegetation plots: **88 %**

Number of habitats with taxon occurrence in the Czech Republic

Number of narrow habitats in which the taxon occurs: **31**
Number of narrow habitats in which the taxon has its optimum: **9**
Number of broad habitats in which the taxon occurs: **9**
Number of broad habitats in which the taxon has its optimum: **5**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz