
Centaurium pulchellum

Distribution





Map info

revised records

unrevised records

On the map are not visualized records without the coordinates and records marked as incorrect or doubtful.





Habitus and growth type

Height [m]: **0.01–0.2**
Growth form: **annual herb**
Life form: **therophyte**
Life strategy: **SR – stress-tolerator/ruderal**
Life strategy (Pierce method based on leaf traits): **R**
Life strategy (Pierce method, C-score): **0 %**
Life strategy (Pierce method, S-score): **13.2 %**
Life strategy (Pierce method, R-score): **86.8 %**

Leaf

Leaf presence and metamorphosis: **leaves present, not modified**
Leaf arrangement (phyllotaxis): **opposite**
Leaf shape: **simple – entire**
Stipules: **absent**
Petiole: **absent**
Leaf life span: **summer green**
Leaf anatomy: **helomorphic**

Flower

Flowering period [month]: **May–July**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Flowering phase: **8 Clematis vitalba-Galium sylvaticum (mid-summer)**
Flower colour: **white, pink**
Flower symmetry: **actinomorphic**
Perianth type: **calyx and corolla**
Perianth fusion: **fused**
Shape of the sympetalous corolla or syntepalous perianth: **hypocrateriform**
Calyx fusion: **synsepalous**
Inflorescence type: **panicula, dichasium**
Dicliny: **synoecious**
Generative reproduction type: **facultative autogamy**
Pollination syndrome: **insect-pollination, selfing**

Fruit, seed and dispersal

Fruit type: **dry fruit – capsule**
Fruit colour: **brown**
Reproduction type: **only by seed/spores**
Dispersal unit (diaspore): **seed**
Dispersal strategy: **Allium (mainly autochory)**
Myrmecochory: **non-myrmecochorous (b)**

Belowground organs and clonality

Shoot life span (cyclicity): **monocyclic shoots prevailing**
Primary root: **present**

Bud bank

Number of buds per shoot at the soil surface (root buds excluded): **5**
Number of buds per shoot at a depth of 0–10 cm (root buds excluded): **0**
Number of buds per shoot at a depth greater than 10 cm (root buds excluded): **0**
Size of the belowground bud bank (root buds excluded): **5**
Depth of the belowground bud bank (root buds exluded) [cm]: **1**
Number of buds per shoot at the soil surface (root buds included): **5**
Number of buds per shoot at a depth of 0–10 cm (root buds included): **0**
Number of buds per shoot at a depth greater than 10 cm (root buds included): **0**
Size of the belowground bud bank (root buds included): **5**
Depth of the belowground bud bank (root buds included) [cm]: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotrophic**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Karyology

Chromosome number (2n): **38 (36, 40, 42, 54)**
Ploidy level (x): **4**
2C genome size [Mbp]: **2350.68**
1Cx monoploid genome size [Mbp]: **587.67**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Genomic GC content: **40.7 %**

Taxon origin

Origin in the Czech Republic: **native**

Ecological indicator values

Ellenberg-type indicator values

Light indicator value: **9 – full light plant, occurring only in fully irradiated places, not at less than 50% of diffuse radiation incident in an open area**
Temperature indicator value: **6 – transition between values 5 and 7**
Moisture indicator value: **7x – humidity indicator, focus on well moistened, but not wet soils (generalist)**
Reaction indicator value: **8 – transition between values 7 and 9, occurring mostly in calcium-rich conditions**
Nutrient indicator value: **4 – transition between values 3 and 5**
Salinity indicator value: **3 – β-mesohaline, mostly on soils with low salt content**

Indicator values for disturbance

Whole-community disturbance frequency indicator value: **-0.51**
Herb layer disturbance frequency indicator value: **-0.09**
Whole-community disturbance severity indicator value: **0.35**
Herb layer disturbance severity indicator value: **0.38**
Whole-community structure based disturbance indicator value: **0.54**
Herb layer structure-based disturbance indicator value: **0.57**

Habitat and sociology

Occurrence in habitats

4 Wetland and riverine herbaceous vegetation

4B Halophilous reed and sedge beds: **1 – rare occurrence**
4H Vegetation of low annual hygrophilous herbs: **1 – rare occurrence**

6 Meadows and mesic pastures

6C Pastures and park grasslands: **1 – rare occurrence**
6G Vegetation of wet disturbed soils: **2 – optimum**

8 Dry grasslands

8D Broad-leaved dry grasslands: **1 – rare occurrence**

10 Saline vegetation

10I Inland saline meadows: **2 – optimum**

13 Anthropogenic vegetation

13C Annual vegetation of trampled habitats: **2 – optimum**

Affinity to the forest environment

Affinity to the forest environment in Thermophyticum: **0 – taxon that does not spontaneously occur in Czech forests**
Affinity to the forest environment in Mesophyticum and Oreophyticum: **0 – taxon that does not spontaneously occur in Czech forests**

Diagnostic taxon

Diagnostic taxon of alliances: **MAC *Verbenion supinae***

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Diagnostic taxon of associations: **MAC01 *Veronico anagalloidis-Lythretum hyssopifoliae*, MCC12 *Tripleurospermo inodori-Bolboschoenetum planiculmis***

Ecological specialization indices

Ecological specialization index for all vegetation types: **4.3**
Ecological specialization index for non-forest vegetation: **4.4**

Colonization ability

Index of colonization success (ICS): **1**
Index of colonization potential (ICP): **1**

Distribution and frequency

Floristic zone: **northern temperate, southern temperate, submeridional, meridional, subtropical**
Floristic region: **Europe, Asia, Africa**
Distribution range extension along the continentality gradient: **10**
Elevational belt in the Czech Republic: **lowlands, colline belt**
Occurrence frequency in the basic grid mapping cells and quadrants of the basic grid mapping cells: 223
taxon.data.freq_in_quad: 423

Commonness in vegetation plots from the Czech Republic

Occurrence frequency in vegetation plots: **0.2 %**
Occurrence frequency in vegetation plots with a cover above 5%: **4.3 %**
Occurrence frequency in vegetation plots with a cover above 25%: **0 %**
Occurrence frequency in vegetation plots with a cover above 50%: **0 %**
Mean percentage cover in vegetation plots: **2.4 %**
Maximum percentage cover in vegetation plots: **13 %**

Number of habitats with taxon occurrence in the Czech Republic

Number of narrow habitats in which the taxon occurs: **7**
Number of narrow habitats in which the taxon has its optimum: **3**
Number of broad habitats in which the taxon occurs: **5**
Number of broad habitats in which the taxon has its optimum: **3**

Threats and protection

Red List 2017 (national categories): **C3 – vulnerable taxon**
Red List 2017 (IUCN categories): **VU – vulnerable**
Legal protection: **not protected by law**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz