
Crambe tataria

Distribution





Map info

revised records

unrevised records

On the map are not visualized records without the coordinates and records marked as incorrect or doubtful.









Habitus and growth type

Height [m]: **0.3–0.8**
Growth form: **monocarpic perennial non-clonal herb**
Life form: **hemicryptophyte**

Leaf

Leaf presence and metamorphosis: **leaves present, not modified**
Leaf arrangement (phyllotaxis): **rosulate**
Leaf shape: **simple – pinnately divided**
Stipules: **absent**
Petiole: **present**

Flower

Flowering period [month]: **May–June**
Flower colour: **white**
Flower symmetry: **actinomorphic**
Perianth type: **calyx and corolla**
Perianth fusion: **free**
Calyx fusion: **aposepalous**
Inflorescence type: **corymbothyrsus**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Fruit, seed and dispersal



Fruit type: **dry fruit – achene/cypsela/samara**
Dispersal unit (diaspore): **fruit, infrutescence or its part, tumbleweed**
Dispersal strategy: **Allium (mainly autochory)**
Myrmecochory: **non-myrmecochorous (b)**



Trophic mode

Parasitism and mycoheterotrophy: **autotrophic**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**



Karyology

Chromosome number (2n): **30 (60, 120)**
Ploidy level (x): **2 (4, 8)**
2C genome size [Mbp]: **2092.97**
1Cx monoploid genome size [Mbp]: **1046.49**
Genomic GC content: **40.5 %**

Taxon origin



Origin in the Czech Republic: **native**

Ecological indicator values



Ellenberg-type indicator values

Light indicator value: **9 – full light plant, occurring only in fully irradiated places, not at less than 50% of diffuse radiation incident in an open area**
Temperature indicator value: **8 – transition between values 7 and 9**
Moisture indicator value: **3 – missing on damp soil**
Reaction indicator value: **8 – transition between values 7 and 9, occurring mostly in calcium-rich conditions**
Nutrient indicator value: **3 – occurring at nutrient-poor sites more frequently than at average sites and exceptionally at rich sites**
Salinity indicator value: **0 – not salt tolerant, glycophyte**



Indicator values for disturbance

Whole-community disturbance frequency indicator value: **0.03**
Herb layer disturbance frequency indicator value: **0.03**
Whole-community disturbance severity indicator value: **0.3**
Herb layer disturbance severity indicator value: **0.34**
Whole-community structure based disturbance indicator value: **0.76**
Herb layer structure-based disturbance indicator value: **0.83**



Habitat and sociology

Occurrence in habitats

8 Dry grasslands

8C Narrow-leaved sub-continental steppes: **2 – optimum**
8D Broad-leaved dry grasslands: **2 – optimum**
8F Thermophilous forest fringe vegetation: **1 – rare occurrence**

13 Anthropogenic vegetation

13D Perennial thermophilous ruderal vegetation: **1 – rare occurrence**

Affinity to the forest environment

Affinity to the forest environment in Thermophyticum: **0 – taxon that does not spontaneously occur in Czech forests**
Affinity to the forest environment in Mesophyticum and Oreophyticum: **0 – taxon that does not spontaneously occur in Czech forests**

Diagnostic taxon

Diagnostic taxon of associations: **THD06 *Astragalo exscapi-Crambetum tatariae***

Constant taxon

Constant taxon of associations: **THD06 *Astragalo exscapi-Crambetum tatariae***

Ecological specialization indices

Ecological specialization index for all vegetation types: **5.9**
Ecological specialization index for non-forest vegetation: **5.8**

Colonization ability

Index of colonization success (ICS): **1**
Index of colonization potential (ICP): **1**

Distribution and frequency

Elevational belt in the Czech Republic: **colline belt**
Occurrence frequency in the basic grid mapping cells and quadrants of the basic grid mapping cells: 17
taxon.data.freq_in_quad: 23

Commonness in vegetation plots from the Czech Republic

Occurrence frequency in vegetation plots: **0.1 %**
Occurrence frequency in vegetation plots with a cover above 5%: **0 %**
Occurrence frequency in vegetation plots with a cover above 25%: **0 %**
Occurrence frequency in vegetation plots with a cover above 50%: **0 %**
Mean percentage cover in vegetation plots: **1.9 %**
Maximum percentage cover in vegetation plots: **3 %**

Number of habitats with taxon occurrence in the Czech Republic

Number of narrow habitats in which the taxon occurs: **4**
Number of narrow habitats in which the taxon has its optimum: **2**
Number of broad habitats in which the taxon occurs: **2**
Number of broad habitats in which the taxon has its optimum: **1**

Threats and protection

Red List 2017 (national categories): **C2r – endangered taxon, rare**
Red List 2017 (IUCN categories): **EN – endangered**
Legal protection: **critically threatened taxon**











© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz