
Dactylorhiza maculata

Distribution





Map info

revised records

unrevised records

On the map are not visualized records without the coordinates and records marked as incorrect or doubtful.





Habitus and growth type

Height [m]: **0.25–0.7**
Growth form: **clonal herb**
Life form: **geophyte**
Life strategy: **CSR – competitor/stress-tolerator/ruderal**

Leaf

Leaf presence and metamorphosis: **leaves present, not modified**
Leaf arrangement (phyllotaxis): **alternate**
Leaf shape: **simple – entire**
Stipules: **absent**
Petiole: **absent**
Leaf life span: **summer green**
Leaf anatomy: **mesomorphic**

Flower

Flowering period [month]: **June–July**
Flowering phase: **6 Cornus sanguinea-Melica uniflora (start of early summer)**
Flower colour: **white, pink-violet, red-violet**
Flower symmetry: **zygomorphic**
Perianth type: **homochlamydeous**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Perianth fusion: **free**
Inflorescence type: **spica**
Dicliny: **synoecious**
Generative reproduction type: **facultative allogamy**
Pollination syndrome: **insect-pollination**



Fruit, seed and dispersal

Fruit type: **dry fruit – capsule**
Fruit colour: **green, brown**
Reproduction type: **only by seed/spores**
Dispersal unit (diaspore): **seed**
Dispersal strategy: **Lycopodium (mainly anemochory)**
Myrmecochory: **non-myrmecochorous (b)**



Belowground organs and clonality

Root metamorphosis: **root tuber**
Storage organ: **root tuber**

Trophic mode

Parasitism and mycoheterotrophy: **partial or initial mycoheterotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**



Taxon origin

Origin in the Czech Republic: **native**

Ecological indicator values

Ellenberg-type indicator values

Light indicator value: **7 – half-light plant, mostly occurring at full light, but also in the shade up to about 30% of diffuse radiation incident in an open area**
Temperature indicator value: **4 – transition between values 3 and 5**
Moisture indicator value: **8 – transition between values 7 and 9**
Reaction indicator value: **5x – indicator of moderate acidity, occurring rarely in strongly acidic as well as in neutral to alkaline conditions (generalist)**
Nutrient indicator value: **3 – occurring at nutrient-poor sites more frequently than at average sites and exceptionally at rich sites**
Salinity indicator value: **0 – not salt tolerant, glycophyte**



Habitat and sociology

Affinity to the forest environment

Affinity to the forest environment in Thermophyticum: **2.1 – taxon occurring both in the forest and open vegetation**
Affinity to the forest environment in Mesophyticum and Oreophyticum: **2.1 – taxon occurring both in the forest and open vegetation**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz









Colonization ability

Index of colonization success (ICS): **1**
Index of colonization potential (ICP): **1**

Distribution and frequency

Floristic zone: **boreal, northern temperate, southern temperate, submeridional**
Floristic region: **Europe**
Continentality degree: **2**
Distribution range extension along the continentality gradient: **2**
Elevational belt in the Czech Republic: **colline belt, submontane belt, montane belt**
Occurrence frequency in the basic grid mapping cells and quadrants of the basic grid mapping cells: 177
taxon.data.freq_in_quad: 266

Threats and protection

Red List 2017 (national categories): **C1b – critically threatened taxon, rare and declining**
Red List 2017 (IUCN categories): **EN – endangered**
Legal protection: **critically threatened taxon**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz







© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz