Dipsacus laciniatus

Distribution



Map info

revised records

unrevised records

On the map are not visualized records without the coordinates and records marked as incorrect or doubtful.

Habitus and growth type

Height [m]: **0.8–2**
Growth form: **monocarpic perennial non-clonal herb**
Life form: **hemicryptophyte**
Life strategy: **CR – competitor/ruderal**

Leaf

Leaf presence and metamorphosis: **leaves present, not modified**
Leaf arrangement (phyllotaxis): **opposite, rosulate**
Leaf shape: **simple – pinnately divided**
Stipules: **absent**
Petiole: **both present and absent**
Leaf anatomy: **scleromorphic**

Flower

Flowering period [month]: **July–September**
Flower colour: **white, pink-violet**
Flower symmetry: **zygomorphic**
Perianth type: **calyx reduced, corolla present**
Perianth fusion: **fused**
Shape of the sympetalous corolla or syntepalous perianth: **tubular**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Calyx fusion: **synsepalous**
Inflorescence type: **capitulum**
Dicliny: **synoecious**
Generative reproduction type: **facultative allogamy**
Pollination syndrome: **insect-pollination**

Fruit, seed and dispersal

Fruit type: **dry fruit – achene/cypsela/samara**
Fruit colour: **brown, grey**
Reproduction type: **only by seed/spores**
Dispersal unit (diaspore): **fruit, infrutescence or its part**
Dispersal strategy: **Allium (mainly autochory)**
Myrmecochory: **probably myrmecochorous**

Belowground organs and clonality

Root metamorphosis: **primary storage root**
Storage organ: **primary storage root**
Shoot life span (cyclicity): **dicyclic or polycyclic shoots prevailing**
Primary root: **present**

Bud bank

Number of buds per shoot at the soil surface (root buds excluded): **5**
Number of buds per shoot at a depth of 0–10 cm (root buds excluded): **0**
Number of buds per shoot at a depth greater than 10 cm (root buds excluded): **0**
Size of the belowground bud bank (root buds excluded): **5**
Depth of the belowground bud bank (root buds exluded) [cm]: **1**
Number of buds per shoot at the soil surface (root buds included): **5**
Number of buds per shoot at a depth of 0–10 cm (root buds included): **0**
Number of buds per shoot at a depth greater than 10 cm (root buds included): **0**
Size of the belowground bud bank (root buds included): **5**
Depth of the belowground bud bank (root buds included) [cm]: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotrophic**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Karyology

Chromosome number (2n): **18 (16)**
Ploidy level (x): **2**
2C genome size [Mbp]: **5834.11**
1Cx monoploid genome size [Mbp]: **2917.05**
Genomic GC content: **38.1 %**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Taxon origin

Origin in the Czech Republic: **native**

Ecological indicator values

Ellenberg-type indicator values

Light indicator value: **8 – light plant, only exceptionally occurring at less than 40% of diffuse radiation incident in an open area**
Temperature indicator value: **7 – heat indicator, occurring in relatively warm lowlands**
Moisture indicator value: **6 – transition between values 5 and 7**
Reaction indicator value: **7 – indicator of slightly acidic to slightly basic conditions, never occurring in very acidic conditions**
Nutrient indicator value: **6 – transition between values 5 and 7**
Salinity indicator value: **1 – salt tolerant, mostly on low-salt to salt-free soils, but occasionally on slightly salty soils**

Habitat and sociology

Occurrence in habitats

4 Wetland and riverine herbaceous vegetation

4A Reed-beds of eutrophic still waters: **1 – rare occurrence**
4B Halophilous reed and sedge beds: **1 – rare occurrence**
4L Nitrophilous herbaceous fringes of lowland rivers: **1 – rare occurrence**

6 Meadows and mesic pastures

6A Mesic Arrhenatherum meadows: **1 – rare occurrence**
6C Pastures and park grasslands: **1 – rare occurrence**
6D Alluvial meadows of lowland rivers: **1 – rare occurrence**

8 Dry grasslands

8F Thermophilous forest fringe vegetation: **2 – optimum**

10 Saline vegetation

10I Inland saline meadows: **1 – rare occurrence**

11 Heathlands and scrub

11J Willow galleries of loamy and sandy river banks: **1 – rare occurrence**
11L Tall mesic and xeric shrub: **2 – optimum**
11R Scrub and pioneer woodland of forests clearings: **1 – rare occurrence**

13 Anthropogenic vegetation

13A Annual vegetation of ruderal habitats: **1 – rare occurrence**
13D Perennial thermophilous ruderal vegetation: **2 – optimum**
13E Perennial nitrophilous herbaceous vegetation of mesic sites: **1 – rare occurrence**
13F Herbaceous vegetation of forests clearings and Rubus scrub: **1 – rare occurrence**

Affinity to the forest environment

Affinity to the forest environment in Thermophyticum: **0 – taxon that does not spontaneously occur in Czech forests**
Affinity to the forest environment in Mesophyticum and Oreophyticum: **0 – taxon that does not spontaneously occur in Czech forests**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Diagnostic taxon

Diagnostic taxon of associations: **KBB05 *Rhamno catharticae-Cornetum sanguineae***

Ecological specialization indices

Ecological specialization index for all vegetation types: **5.3**
Ecological specialization index for non-forest vegetation: **5.3**

Colonization ability

Index of colonization success (ICS): **1**

Distribution and frequency

Floristic zone: **southern temperate, submeridional, meridional**
Floristic region: **Europe, Western Asia**
Continentality degree: **7**
Distribution range extension along the continentality gradient: **5**
Elevational belt in the Czech Republic: **lowlands, colline belt**
Occurrence frequency in the basic grid mapping cells and quadrants of the basic grid mapping cells:
taxon.data.freq_in_quad:

Commonness in vegetation plots from the Czech Republic

Occurrence frequency in vegetation plots: **0 %**
Occurrence frequency in vegetation plots with a cover above 5%: **0 %**
Occurrence frequency in vegetation plots with a cover above 25%: **0 %**
Occurrence frequency in vegetation plots with a cover above 50%: **0 %**
Mean percentage cover in vegetation plots: **2.3 %**
Maximum percentage cover in vegetation plots: **3 %**

Number of habitats with taxon occurrence in the Czech Republic

Number of narrow habitats in which the taxon occurs: **15**
Number of narrow habitats in which the taxon has its optimum: **3**
Number of broad habitats in which the taxon occurs: **6**
Number of broad habitats in which the taxon has its optimum: **3**

Threats and protection

Red List 2017 (national categories): **C3 – vulnerable taxon**
Red List 2017 (IUCN categories): **NT – near threatened**
Legal protection: **not protected by law**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz