
Epipactis greuteri

Distribution





Map info

revised records

unrevised records

On the map are not visualized records without the coordinates and records marked as incorrect or doubtful.





Habitus and growth type

Height [m]: **0.3–0.65**
Growth form: **clonal herb**
Life form: **geophyte**

Leaf

Leaf presence and metamorphosis: **leaves present, not modified**
Leaf arrangement (phyllotaxis): **alternate**
Leaf shape: **simple – entire**
Stipules: **absent**
Petiole: **absent**

Flower

Flowering period [month]: **July–August**
Flower colour: **green-white, green**
Flower symmetry: **zygomorphic**
Perianth type: **homochlamydeous**
Perianth fusion: **free**
Inflorescence type: **spica**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz







Fruit, seed and dispersal

Fruit type: **dry fruit – capsule**
Dispersal unit (diaspore): **seed**
Dispersal strategy: **Lycopodium (mainly anemochory)**
Myrmecochory: **non-myrmecochorous (b)**

Trophic mode

Parasitism and mycoheterotrophy: **partial or initial mycoheterotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in the Czech Republic: **native**

Ecological indicator values

Ellenberg-type indicator values

Light indicator value: **2 – transition between values 1 and 3**
Temperature indicator value: **6 – transition between values 5 and 7**
Moisture indicator value: **7 – humidity indicator, focus on well moistened, but not wet soils**
Reaction indicator value: **7 – indicator of slightly acidic to slightly basic conditions, never occurring in very acidic conditions**
Nutrient indicator value: **6 – transition between values 5 and 7**
Salinity indicator value: **0 – not salt tolerant, glycophyte**

Habitat and sociology

Occurrence in habitats

12 Forests

12C Oak-hornbeam forests: **1 – rare occurrence**
12D Ravine forests: **1 – rare occurrence**
12E Herb-rich beech forests: **2 – optimum**
12F Limestone beech forests: **1 – rare occurrence**
12G Acidophilous beech forests: **1 – rare occurrence**
12V Spruce plantations: **2 – optimum**

Affinity to the forest environment

Affinity to the forest environment in Thermophyticum: **1.1 – taxon occurring mainly in the closed forest**
Affinity to the forest environment in Mesophyticum and Oreophyticum: **1.1 – taxon occurring mainly in the closed forest**

Distribution and frequency

Elevational belt in the Czech Republic: **colline belt, submontane belt**
Occurrence frequency in the basic grid mapping cells and quadrants of the basic grid

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

mapping cells: 36
taxon.data.freq_in_quad: 49

Number of habitats with taxon occurrence in the Czech Republic

Number of narrow habitats in which the taxon occurs: **6**
Number of narrow habitats in which the taxon has its optimum: **2**
Number of broad habitats in which the taxon occurs: **1**
Number of broad habitats in which the taxon has its optimum: **1**

Threats and protection

Red List 2017 (national categories): **C2r – endangered taxon, rare**
Red List 2017 (IUCN categories): **VU – vulnerable**
Legal protection: **not protected by law**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz