
Epipactis leptochila

Distribution






Map info

revised records

unrevised records

On the map are not visualized records without the coordinates and records marked as incorrect or doubtful.







Habitus and growth type

Height [m]: **0.25–0.7**
Growth form: **clonal herb**
Life form: **geophyte**
Life strategy: **CSR – competitor/stress-tolerator/ruderal**

Leaf

Leaf presence and metamorphosis: **leaves present, not modified**
Leaf arrangement (phyllotaxis): **alternate**
Leaf shape: **simple – entire**
Stipules: **absent**
Petiole: **absent**
Leaf life span: **summer green**
Leaf anatomy: **mesomorphic, hygromorphic**

Flower

Flowering period [month]: **July**
Flowering phase: **8 Clematis vitalba-Galium sylvaticum (mid-summer)**
Flower colour: **green**
Flower symmetry: **zygomorphic**
Perianth type: **homochlamydeous**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Perianth fusion: **free**
Inflorescence type: **spica**
Dicliny: **synoecious**
Generative reproduction type: **autogamy**
Pollination syndrome: **insect-pollination, selfing, cleistogamy**

Fruit, seed and dispersal

Fruit type: **dry fruit – capsule**
Reproduction type: **by seed/spores and vegetatively**
Dispersal unit (diaspore): **seed**
Dispersal strategy: **Lycopodium (mainly anemochory)**
Myrmecochory: **non-myrmecochorous (b)**

Belowground organs and clonality

Shoot metamorphosis: **rhizome**
Storage organ: **rhizome**
Type of clonal growth organ: **hypogeogenous rhizome**
Freely dispersible organs of clonal growth: **absent**
Shoot life span (cyclicity): **monocyclic shoots prevailing**
Branching type of stem-derived organs of clonal growth: **sympodial**
Primary root: **absent**
Persistence of the clonal growth organ [year]: **4**
Number of clonal offspring: **1**
Lateral spreading distance by clonal growth [m]: **0.01**
Clonal index: **3**

Bud bank

Number of buds per shoot at the soil surface (root buds excluded): **5**
Number of buds per shoot at a depth of 0–10 cm (root buds excluded): **15**
Number of buds per shoot at a depth greater than 10 cm (root buds excluded): **0**
Size of the belowground bud bank (root buds excluded): **20**
Depth of the belowground bud bank (root buds exluded) [cm]: **4**
Number of buds per shoot at the soil surface (root buds included): **5**
Number of buds per shoot at a depth of 0–10 cm (root buds included): **15**
Number of buds per shoot at a depth greater than 10 cm (root buds included): **0**
Size of the belowground bud bank (root buds included): **20**
Depth of the belowground bud bank (root buds included) [cm]: **4**

Trophic mode

Parasitism and mycoheterotrophy: **partial or initial mycoheterotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in the Czech Republic: **native**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Ecological indicator values

Ellenberg-type indicator values

Light indicator value: **3 – shade plant, usually occurring where the incident radiation is less than 5% of that in an open area, but also at sunnier sites**
Temperature indicator value: **6 – transition between values 5 and 7**
Moisture indicator value: **4 – transition between values 3 and 5**
Reaction indicator value: **7 – indicator of slightly acidic to slightly basic conditions, never occurring in very acidic conditions**
Nutrient indicator value: **6 – transition between values 5 and 7**
Salinity indicator value: **0 – not salt tolerant, glycophyte**

Habitat and sociology

Occurrence in habitats

12 Forests

12C Oak-hornbeam forests: **1 – rare occurrence**
12E Herb-rich beech forests: **1 – rare occurrence**

Affinity to the forest environment

Affinity to the forest environment in Thermophyticum: **1.1 – taxon occurring mainly in the closed forest**
Affinity to the forest environment in Mesophyticum and Oreophyticum: **1.1 – taxon occurring mainly in the closed forest**

Distribution and frequency

Floristic zone: **northern temperate, southern temperate**
Floristic region: **Europe**
Continentality degree: **4**
Distribution range extension along the continentality gradient: **4**
Elevational belt in the Czech Republic: **colline belt, submontane belt**
Occurrence frequency in the basic grid mapping cells and quadrants of the basic grid mapping cells: 12
taxon.data.freq_in_quad: 13

Number of habitats with taxon occurrence in the Czech Republic

Number of narrow habitats in which the taxon occurs: **2**
Number of narrow habitats in which the taxon has its optimum: **0**
Number of broad habitats in which the taxon occurs: **1**
Number of broad habitats in which the taxon has its optimum: **0**

Threats and protection

Red List 2017 (national categories): **C1r – critically threatened taxon, rare**
Red List 2017 (IUCN categories): **CR – critically endangered**
Legal protection: **critically threatened taxon**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz