


# *Epipactis leptochila*

## Distribution



### Map info

● revised records

○ unrevised records

On the map are not visualized records without the coordinates and records marked as incorrect or doubtful.

## Habitus and growth type

Height [m]: **0.25-0.7**

Growth form: **clonal herb**

Life form: **geophyte**

Life strategy: **CSR - competitor/stress-tolerator/ruderal**

## Leaf

Leaf presence and metamorphosis: **leaves present, not modified**

Leaf arrangement (phyllotaxis): **alternate**

Leaf shape: **simple - entire**

Stipules: **absent**

Petiole: **absent**

Leaf life span: **summer green**

Leaf anatomy: **mesomorphic, hygromorphic**

## Flower

Flowering period [month]: **July**

Flowering phase: **8 Clematis vitalba-Galium sylvaticum (mid-summer)**

Flower colour: **green**

Flower symmetry: **zygomorphic**

Perianth type: **homochlamydeous**



Perianth fusion: **free**  
Inflorescence type: **spica**  
Dicliny: **synoecious**  
Generative reproduction type: **autogamy**  
Pollination syndrome: **insect-pollination, selfing, cleistogamy**

### **Fruit, seed and dispersal**

Fruit type: **dry fruit - capsule**  
Reproduction type: **by seed/spores and vegetatively**  
Dispersal unit (diaspore): **seed**  
Dispersal strategy: **Lycopodium (mainly anemochory)**  
Myrmecochory: **non-myrmecochorous (b)**

### **Belowground organs and clonality**

Shoot metamorphosis: **rhizome**  
Storage organ: **rhizome**  
Type of clonal growth organ: **hypogeogenous rhizome**  
Freely dispersible organs of clonal growth: **absent**  
Shoot life span (cyclicality): **monocyclic shoots prevailing**  
Branching type of stem-derived organs of clonal growth: **sympodial**  
Primary root: **absent**  
Persistence of the clonal growth organ [year]: **4**  
Number of clonal offspring: **1**  
Lateral spreading distance by clonal growth [m]: **0.01**  
Clonal index: **3**

### **Bud bank**

Number of buds per shoot at the soil surface (root buds excluded): **5**  
Number of buds per shoot at a depth of 0–10 cm (root buds excluded): **15**  
Number of buds per shoot at a depth greater than 10 cm (root buds excluded): **0**  
Size of the belowground bud bank (root buds excluded): **20**  
Depth of the belowground bud bank (root buds excluded) [cm]: **4**  
Number of buds per shoot at the soil surface (root buds included): **5**  
Number of buds per shoot at a depth of 0–10 cm (root buds included): **15**  
Number of buds per shoot at a depth greater than 10 cm (root buds included): **0**  
Size of the belowground bud bank (root buds included): **20**  
Depth of the belowground bud bank (root buds included) [cm]: **4**

### **Trophic mode**

Parasitism and mycoheterotrophy: **partial or initial mycoheterotroph**  
Carnivory: **non-carnivorous**  
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

### **Taxon origin**

Origin in the Czech Republic: **native**

## Ecological indicator values

### Ellenberg-type indicator values

Light indicator value: **3 - shade plant, usually occurring where the incident radiation is less than 5% of that in an open area, but also at sunnier sites**

Temperature indicator value: **6 - transition between values 5 and 7**

Moisture indicator value: **4 - transition between values 3 and 5**

Reaction indicator value: **7 - indicator of slightly acidic to slightly basic conditions, never occurring in very acidic conditions**

Nutrient indicator value: **6 - transition between values 5 and 7**

Salinity indicator value: **0 - not salt tolerant, glycophyte**

## Habitat and sociology

### Occurrence in habitats

12 Forests

12C Oak-hornbeam forests: **1 - rare occurrence**

12E Herb-rich beech forests: **1 - rare occurrence**

### Affinity to the forest environment

Affinity to the forest environment in Thermophyticum: **1.1 - taxon occurring mainly in the closed forest**

Affinity to the forest environment in Mesophyticum and Oreophyticum: **1.1 - taxon occurring mainly in the closed forest**

## Distribution and frequency

Floristic zone: **northern temperate, southern temperate**

Floristic region: **Europe**

Continental degree: **4**

Distribution range extension along the continentality gradient: **4**

Elevational belt in the Czech Republic: **colline belt, submontane belt**

Occurrence frequency in the basic grid mapping cells and quadrants of the basic grid mapping cells: **12**

taxon.data.freq\_in\_quad: **13**

Number of habitats with taxon occurrence in the Czech Republic

Number of narrow habitats in which the taxon occurs: **2**

Number of narrow habitats in which the taxon has its optimum: **0**

Number of broad habitats in which the taxon occurs: **1**

Number of broad habitats in which the taxon has its optimum: **0**

## Threats and protection

Red List 2017 (national categories): **C1r - critically threatened taxon, rare**

Red List 2017 (IUCN categories): **CR - critically endangered**

Legal protection: **critically threatened taxon**