
Galeopsis pubescens

Distribution





Map info

revised records

unrevised records

On the map are not visualized records without the coordinates and records marked as incorrect or doubtful.





Habitus and growth type

Height [m]: **0.2–0.5**
Growth form: **annual herb**
Life form: **therophyte**
Life strategy: **CR – competitor/ruderal**

Leaf

Leaf presence and metamorphosis: **leaves present, not modified**
Leaf arrangement (phyllotaxis): **opposite**
Leaf shape: **simple – entire**
Stipules: **absent**
Petiole: **present**
Leaf life span: **summer green**
Leaf anatomy: **mesomorphic**

Flower

Flowering period [month]: **July–October**
Flowering phase: **8 Clematis vitalba-Galium sylvaticum (mid-summer)**
Flower colour: **red-violet**
Flower symmetry: **zygomorphic**
Perianth type: **calyx and corolla**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Perianth fusion: **fused**
Shape of the sympetalous corolla or syntepalous perianth: **bilabiate**
Calyx fusion: **synsepalous**
Inflorescence type: **pseudospica e verticillastris composita**
Dicliny: **synoecious, gynomonoecious**
Generative reproduction type: **facultative allogamy**
Pollination syndrome: **insect-pollination**



Fruit, seed and dispersal

Fruit type: **dry fruit – cluster of four one-seeded nutlets**
Fruit colour: **brown, black**
Reproduction type: **only by seed/spores**
Dispersal unit (diaspore): **fruit, infrutescence or its part**
Dispersal strategy: **Allium (mainly autochory)**
Myrmecochory: **non-myrmecochorous (b) nv**



Belowground organs and clonality

Shoot life span (cyclicity): **monocyclic shoots prevailing**
Primary root: **present**

Bud bank

Number of buds per shoot at the soil surface (root buds excluded): **5**
Number of buds per shoot at a depth of 0–10 cm (root buds excluded): **0**
Number of buds per shoot at a depth greater than 10 cm (root buds excluded): **0**
Size of the belowground bud bank (root buds excluded): **5**
Depth of the belowground bud bank (root buds exluded) [cm]: **1**
Number of buds per shoot at the soil surface (root buds included): **5**
Number of buds per shoot at a depth of 0–10 cm (root buds included): **0**
Number of buds per shoot at a depth greater than 10 cm (root buds included): **0**
Size of the belowground bud bank (root buds included): **5**
Depth of the belowground bud bank (root buds included) [cm]: **1**





Trophic mode

Parasitism and mycoheterotrophy: **autotrophic**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**



Karyology

Chromosome number (2n): **16**
Ploidy level (x): **2**
2C genome size [Mbp]: **1614.14**
1Cx monoploid genome size [Mbp]: **807.07**
Genomic GC content: **38.7 %**



© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Taxon origin



Origin in the Czech Republic: **native**

Ecological indicator values

Ellenberg-type indicator values

Light indicator value: **6 – transition between values 5 and 7; rarely at less than 20% of diffuse radiation incident in an open area**
Temperature indicator value: **5 – moderate heat indicator, occurring from lowland to montane belt, mainly in submontane-temperate areas**
Moisture indicator value: **5 – indicator of fresh soils, focus on soils of average moisture, missing on wet and on soils that frequently dry out**
Reaction indicator value: **6x – transition between values 5 and 7 (generalist)**
Nutrient indicator value: **7 – occurring at nutrient-rich sites more often than at average sites and only exceptionally at poor sites**
Salinity indicator value: **0 – not salt tolerant, glycophyte**



Indicator values for disturbance

Whole-community disturbance frequency indicator value: **-1.7**
Herb layer disturbance frequency indicator value: **-0.73**
Whole-community disturbance severity indicator value: **0.45**
Herb layer disturbance severity indicator value: **0.36**
Whole-community structure based disturbance indicator value: **0.27**
Herb layer structure-based disturbance indicator value: **0.34**

Habitat and sociology

Occurrence in habitats

1 Vegetation of cliffs, screes and walls

1D Mobile calcareous screes: **1 – rare occurrence**

2 Alpine and subalpine grasslands

2B Subalpine tall-forb and tall-grass vegetation: **1 – rare occurrence**

4 Wetland and riverine herbaceous vegetation

4D Riverine reed vegetation: **1 – rare occurrence**
4E Reed vegetation of brooks: **1 – rare occurrence**
4I Vegetation of nitrophilous annual hygrophilous herbs: **1 – rare occurrence**
4K Petasites fringes of montane brooks: **1 – rare occurrence**
4L Nitrophilous herbaceous fringes of lowland rivers: **1 – rare occurrence**

6 Meadows and mesic pastures

6E Wet Cirsium meadows: **1 – rare occurrence**

8 Dry grasslands

8A Hercynian dry grasslands on rock outcrops: **1 – rare occurrence**
8F Thermophilous forest fringe vegetation: **1 – rare occurrence**

11 Heathlands and scrub

11I Willow carrs: **1 – rare occurrence**
11J Willow galleries of loamy and sandy river banks: **1 – rare occurrence**
11L Tall mesic and xeric shrub: **1 – rare occurrence**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

11R Scrub and pioneer woodland of forests clearings: **2 – optimum**

12 Forests

12B Alluvial forests: **2 – optimum**
12C Oak-hornbeam forests: **2 – optimum**
12D Ravine forests: **2 – optimum**
12E Herb-rich beech forests: **2 – optimum**
12F Limestone beech forests: **1 – rare occurrence**
12G Acidophilous beech forests: **1 – rare occurrence**
12H Peri-Alpidic basiphilous thermophilous oak forests: **1 – rare occurrence**
12I Sub-continental thermophilous oak forests: **1 – rare occurrence**
12J Acidophilous thermophilous oak forests: **2 – optimum**
12K Acidophilous oak forests: **1 – rare occurrence**
12L Boreo-continental pine forests: **1 – rare occurrence**
12T Robinia pseudacacia plantations: **2 – optimum**
12U Plantations of broad-leaved non-native trees: **1 – rare occurrence**
12V Spruce plantations: **2 – optimum**
12W Pine and larch plantations: **1 – rare occurrence**

13 Anthropogenic vegetation

13A Annual vegetation of ruderal habitats: **1 – rare occurrence**
13B Annual vegetation of arable land: **2 – optimum**
13D Perennial thermophilous ruderal vegetation: **1 – rare occurrence**
13E Perennial nitrophilous herbaceous vegetation of mesic sites: **2 – optimum**
13F Herbaceous vegetation of forests clearings and Rubus scrub: **1 – rare occurrence**

Affinity to the forest environment

Affinity to the forest environment in Thermophyticum: **1.2 – taxon occurring mainly along forest edges and in forest openings, including forest roads and paths, windthrow sites, burnt sites and forest clearings**
Affinity to the forest environment in Mesophyticum and Oreophyticum: **1.2 – taxon occurring mainly along forest edges and in forest openings, including forest roads and paths, windthrow sites, burnt sites and forest clearings**

Constant taxon

Constant taxon of associations: **XDC05 *Urtico dioicae-Parietarietum officinalis***

Ecological specialization indices

Ecological specialization index for all vegetation types: **3.7**
Ecological specialization index for non-forest vegetation: **3.4**
Ecological specialization index for forest vegetation: **4.6**

Colonization ability

Index of colonization success (ICS): **6**
Index of colonization potential (ICP): **8**
Optimum successional age [years]: **30**

Distribution and frequency

Floristic zone: **northern temperate, southern temperate, submeridional**
Floristic region: **Europe**
Continentality degree: **6**
Distribution range extension along the continentality gradient: **2**
Elevational belt in the Czech Republic: **lowlands, colline belt, submontane belt,**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

montane belt

Occurrence frequency in the basic grid mapping cells and quadrants of the basic grid mapping cells: 639

taxon.data.freq_in_quad: 2026

Commonness in vegetation plots from the Czech Republic

Occurrence frequency in vegetation plots: **1.3 %**

Occurrence frequency in vegetation plots with a cover above 5%: **4.4 %**

Occurrence frequency in vegetation plots with a cover above 25%: **0.8 %**

Occurrence frequency in vegetation plots with a cover above 50%: **0 %**

Mean percentage cover in vegetation plots: **2.6 %**

Maximum percentage cover in vegetation plots: **38 %**

Number of habitats with taxon occurrence in the Czech Republic

Number of narrow habitats in which the taxon occurs: **34**

Number of narrow habitats in which the taxon has its optimum: **10**

Number of broad habitats in which the taxon occurs: **8**

Number of broad habitats in which the taxon has its optimum: **3**

Threats and protection

Red List 2017 (national categories): **taxon is not on the Red List**

Red List 2017 (IUCN categories): **LC(NA) – least concern (taxon is not on the Red List)**

Legal protection: **not protected by law**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz