
Geranium sylvaticum

Distribution





Map info

revised records

unrevised records

On the map are not visualized records without the coordinates and records marked as incorrect or doubtful.







Habitus and growth type

Height [m]: **0.3–0.6**
Growth form: **clonal herb**
Life form: **hemicryptophyte**
Life strategy: **C – competitor**
Life strategy (Pierce method based on leaf traits): **CR/CSR**
Life strategy (Pierce method, C-score): **36.3 %**
Life strategy (Pierce method, S-score): **24.2 %**
Life strategy (Pierce method, R-score): **39.5 %**

Leaf

Leaf presence and metamorphosis: **leaves present, not modified**
Leaf arrangement (phyllotaxis): **opposite**
Leaf shape: **simple – palmately divided**
Stipules: **present**
Petiole: **present**
Leaf life span: **summer green**
Leaf anatomy: **mesomorphic, hygromorphic**


Flower

Flowering period [month]: **June–August**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz



Flowering phase: **5 Sorbus aucuparia-Galium odoratum (end of mid-spring)**
Flower colour: **pink, violet**
Flower symmetry: **actinomorphic**
Perianth type: **calyx and corolla**
Perianth fusion: **free**
Calyx fusion: **aposepalous**
Inflorescence type: **dichasium**
Dicliny: **synoecious, gynomonoecious, gynodioecious**
Generative reproduction type: **allogamy**
Pollination syndrome: **insect-pollination**
Pollinator spectrum: **solitary bees, flies s. l. (honeybee, bumblebees, other Hymenoptera, hoverflies, meat flies s. l., other Diptera, butterflies, beetles, nitidulids, other pollinators, unknown)**



Fruit, seed and dispersal

Fruit type: **dry fruit – dry schizocarp with an apical beak**
Reproduction type: **by seed/spores and vegetatively**
Dispersal unit (diaspore): **seed, fruit, infrutescence or its part**
Dispersal strategy: **Allium (mainly autochory)**
Myrmecochory: **non-myrmecochorous (b)**

Belowground organs and clonality

Shoot metamorphosis: **rhizome**
Storage organ: **rhizome**
Type of clonal growth organ: **epigeogenous rhizome**
Freely dispersible organs of clonal growth: **absent**
Shoot life span (cyclicity): **dicyclic or polycyclic shoots prevailing**
Branching type of stem-derived organs of clonal growth: **sympodial**
Primary root: **absent**
Persistence of the clonal growth organ [year]: **4**
Number of clonal offspring: **0.9**
Lateral spreading distance by clonal growth [m]: **0.02**
Clonal index: **3**

Bud bank

Number of buds per shoot at the soil surface (root buds excluded): **5**
Number of buds per shoot at a depth of 0–10 cm (root buds excluded): **14**
Number of buds per shoot at a depth greater than 10 cm (root buds excluded): **0**
Size of the belowground bud bank (root buds excluded): **19**
Depth of the belowground bud bank (root buds exluded) [cm]: **4**
Number of buds per shoot at the soil surface (root buds included): **5**
Number of buds per shoot at a depth of 0–10 cm (root buds included): **14**
Number of buds per shoot at a depth greater than 10 cm (root buds included): **0**
Size of the belowground bud bank (root buds included): **19**
Depth of the belowground bud bank (root buds included) [cm]: **4**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Trophic mode

Parasitism and mycoheterotrophy: **autotrophic**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Karyology

Chromosome number (2n): **28**
Ploidy level (x): **2**
2C genome size [Mbp]: **3962.7**
1Cx monoploid genome size [Mbp]: **1981.35**
Genomic GC content: **44.7 %**

Taxon origin

Origin in the Czech Republic: **native**

Ecological indicator values

Ellenberg-type indicator values

Light indicator value: **7 – half-light plant, mostly occurring at full light, but also in the shade up to about 30% of diffuse radiation incident in an open area**
Temperature indicator value: **4 – transition between values 3 and 5**
Moisture indicator value: **6 – transition between values 5 and 7**
Reaction indicator value: **6 – transition between values 5 and 7**
Nutrient indicator value: **6 – transition between values 5 and 7**
Salinity indicator value: **0 – not salt tolerant, glycophyte**

Indicator values for disturbance

Whole-community disturbance frequency indicator value: **-0.93**
Herb layer disturbance frequency indicator value: **-0.43**
Whole-community disturbance severity indicator value: **0.28**
Herb layer disturbance severity indicator value: **0.3**
Whole-community structure based disturbance indicator value: **0.44**
Herb layer structure-based disturbance indicator value: **0.59**

Habitat and sociology

Occurrence in habitats

1 Vegetation of cliffs, screes and walls

1B Siliceous cliffs and block fields: **1 – rare occurrence**

2 Alpine and subalpine grasslands

2B Subalpine tall-forb and tall-grass vegetation: **2 – optimum**

4 Wetland and riverine herbaceous vegetation

4K Petasites fringes of montane brooks: **1 – rare occurrence**

5 Vegetation of springs and mires

5B Lowland to montane soft-water springs: **1 – rare occurrence**
5C Alpine and subalpine soft-water springs: **1 – rare occurrence**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

6 Meadows and mesic pastures

6A Mesic Arrhenatherum meadows: **1 – rare occurrence**

6B Montane mesic meadows: **3 – dominant**

6E Wet Cirsium meadows: **2 – optimum**

7 Acidophilous grasslands

7A Subalpine and montane acidophilous grasslands: **2 – optimum**

7B Submontane Nardus grasslands: **1 – rare occurrence**

11 Heathlands and scrub

11D Subalpine acidophilous Pinus mugo scrub: **1 – rare occurrence**

11H Subalpine deciduous scrub: **2 – optimum**

12 Forests

12B Alluvial forests: **1 – rare occurrence**

13 Anthropogenic vegetation

13E Perennial nitrophilous herbaceous vegetation of mesic sites: **1 – rare occurrence**

Affinity to the forest environment

Affinity to the forest environment in Thermophyticum: **2.2 – taxon occurring partly in the forest, but mainly in open vegetation**

Affinity to the forest environment in Mesophyticum and Oreophyticum: **2.2 – taxon occurring partly in the forest, but mainly in open vegetation**

Diagnostic taxon

Diagnostic taxon of alliances: **ADB *Calamagrostion arundinaceae***, **ADC *Salicion silesiacae***, **ADD *Adenostylion alliariae***, **TDB *Polygono bistortae-Trisetion flavescentis***, **TEB *Nardo strictae-Agrostion tenuis***, **XDF *Rumicion alpini***

Diagnostic taxon of associations: **ADB01 *Bupleuro longifoliae-Calamagrostietum arundinaceae***, **ADC01 *Salici silesiacae-Betuletum carpaticae***, **ADD03 *Trollio altissimi-Geranietum sylvatici***, **ADD04 *Laserpitio archangelicae-Dactylidetum glomeratae***, **TDB01 *Geranio sylvatici-Trisetetum flavescentis***, **TDB02 *Melandrio rubri-Phleetum alpini***, **TEB01 *Sileno vulgaris-Nardetum strictae***, **XDF01 *Rumicetum alpini***

Constant taxon

Constant taxon of alliances: **ADB *Calamagrostion arundinaceae***, **TDB *Polygono bistortae-Trisetion flavescentis***, **TEB *Nardo strictae-Agrostion tenuis***

Constant taxon of associations: **ADB01 *Bupleuro longifoliae-Calamagrostietum arundinaceae***, **ADD03 *Trollio altissimi-Geranietum sylvatici***, **ADD04 *Laserpitio archangelicae-Dactylidetum glomeratae***, **ADE01 *Daphno mezerei-Dryopteridetum filicis-maris***, **KCA02 *Adenostylo alliariae-Pinetum mugo***, **TDB01 *Geranio sylvatici-Trisetetum flavescentis***, **TDB02 *Melandrio rubri-Phleetum alpini***, **TEB01 *Sileno vulgaris-Nardetum strictae***

Dominant taxon

Dominant taxon of associations: **ADD03 *Trollio altissimi-Geranietum sylvatici***, **TDB01 *Geranio sylvatici-Trisetetum flavescentis***, **TDB02 *Melandrio rubri-Phleetum alpini***, **TDB03 *Meo athamantici-Festucetum rubrae***

Ecological specialization indices

Ecological specialization index for all vegetation types: **5.5**

Ecological specialization index for non-forest vegetation: **5.8**

Ecological specialization index for forest vegetation: **5.5**

Colonization ability

Index of colonization success (ICS): **2**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Index of colonization potential (ICP): **1**
Optimum successional age [years]: **8**

Distribution and frequency

Floristic zone: **boreal, northern temperate, southern temperate, submeridional**
Floristic region: **Europe, Western Siberia**
Continentality degree: **6**
Distribution range extension along the continentality gradient: **5**
Elevational belt in the Czech Republic: **submontane belt, montane belt**
Occurrence frequency in the basic grid mapping cells and quadrants of the basic grid mapping cells: 252
taxon.data.freq_in_quad: 490

Commonness in vegetation plots from the Czech Republic

Occurrence frequency in vegetation plots: **1.1 %**
Occurrence frequency in vegetation plots with a cover above 5%: **28.3 %**
Occurrence frequency in vegetation plots with a cover above 25%: **8.5 %**
Occurrence frequency in vegetation plots with a cover above 50%: **0.6 %**
Mean percentage cover in vegetation plots: **7.5 %**
Maximum percentage cover in vegetation plots: **63 %**

Number of habitats with taxon occurrence in the Czech Republic

Number of narrow habitats in which the taxon occurs: **14**
Number of narrow habitats in which the taxon has its optimum: **5**
Number of broad habitats in which the taxon occurs: **9**
Number of broad habitats in which the taxon has its optimum: **4**

Threats and protection

Red List 2017 (national categories): **taxon is not on the Red List**
Red List 2017 (IUCN categories): **LC(NA) – least concern (taxon is not on the Red List)**
Legal protection: **not protected by law**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz