
Hordeum jubatum



Distribution



Map info

revised records

unrevised records

On the map are not visualized records without the coordinates and records marked as incorrect or doubtful.







Habitus and growth type

Height [m]: **0.15–0.4**
Growth form: **annual herb**
Life form: **hemicryptophyte**
Life strategy: **SR – stress-tolerator/ruderal**
Life strategy (Pierce method based on leaf traits): **SR**
Life strategy (Pierce method, C-score): **9.9 %**
Life strategy (Pierce method, S-score): **44.2 %**
Life strategy (Pierce method, R-score): **45.9 %**

Leaf

Leaf presence and metamorphosis: **leaves present, not modified**
Leaf arrangement (phyllotaxis): **alternate**
Leaf shape: **simple – entire**
Stipules: **absent**
Petiole: **absent**
Leaf life span: **summer green**
Leaf anatomy: **mesomorphic, helomorphic**

Flower

Flowering period [month]: **June–September**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz




Flowering phase: **7 Ligustrum vulgare-Stachys sylvatica (end of early summer)**
Flower colour: **green**
Perianth type: **reduced**
Perianth fusion: **reduced**
Inflorescence type: **spica e spiculis composita**
Dicliny: **andromonoecious**
Generative reproduction type: **mixed mating**
Pollination syndrome: **selfing**

Fruit, seed and dispersal



Fruit type: **dry fruit – caryopsis**
Fruit colour: **brown**
Reproduction type: **mostly by seed/spores, rarely vegetatively**
Dispersal unit (diaspore): **fruit, infrutescence or its part**
Dispersal strategy: **Allium (mainly autochory)**
Myrmecochory: **non-myrmecochorous (b)**

Belowground organs and clonality



Storage organ: **tuft**
Shoot life span (cyclicity): **monocyclic shoots prevailing**
Primary root: **absent**

Bud bank

Number of buds per shoot at the soil surface (root buds excluded): **5**
Number of buds per shoot at a depth of 0–10 cm (root buds excluded): **3**
Number of buds per shoot at a depth greater than 10 cm (root buds excluded): **0**
Size of the belowground bud bank (root buds excluded): **8**
Depth of the belowground bud bank (root buds exluded) [cm]: **2**
Number of buds per shoot at the soil surface (root buds included): **5**
Number of buds per shoot at a depth of 0–10 cm (root buds included): **3**
Number of buds per shoot at a depth greater than 10 cm (root buds included): **0**
Size of the belowground bud bank (root buds included): **8**
Depth of the belowground bud bank (root buds included) [cm]: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotrophic**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Karyology

Chromosome number (2n): **28**
Ploidy level (x): **4**
2C genome size [Mbp]: **16186.36**
1Cx monoploid genome size [Mbp]: **4046.59**
Genomic GC content: **47.5 %**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Taxon origin

Origin in the Czech Republic: **neophyte**
Invasion status: **naturalized**
Geographic origin: **North America**
Year of the first record in the wild: **1880**
Period of introduction: **Late Modern Period (1800–1950)**
Introduction pathway: **intentional – ornamental**

Ecological indicator values

Ellenberg-type indicator values

Light indicator value: **9 – full light plant, occurring only in fully irradiated places, not at less than 50% of diffuse radiation incident in an open area**
Temperature indicator value: **6 – transition between values 5 and 7**
Moisture indicator value: **5 – indicator of fresh soils, focus on soils of average moisture, missing on wet and on soils that frequently dry out**
Reaction indicator value: **7 – indicator of slightly acidic to slightly basic conditions, never occurring in very acidic conditions**
Nutrient indicator value: **6 – transition between values 5 and 7**
Salinity indicator value: **3 – β-mesohaline, mostly on soils with low salt content**

Habitat and sociology

Occurrence in habitats

6 Meadows and mesic pastures

6G Vegetation of wet disturbed soils: **2 – optimum**

10 Saline vegetation

10I Inland saline meadows: **2 – optimum**

13 Anthropogenic vegetation

13A Annual vegetation of ruderal habitats: **2 – optimum**
13B Annual vegetation of arable land: **1 – rare occurrence**
13C Annual vegetation of trampled habitats: **1 – rare occurrence**
13D Perennial thermophilous ruderal vegetation: **1 – rare occurrence**

Affinity to the forest environment

Affinity to the forest environment in Thermophyticum: **0 – taxon that does not spontaneously occur in Czech forests**
Affinity to the forest environment in Mesophyticum and Oreophyticum: **0 – taxon that does not spontaneously occur in Czech forests**

Colonization ability

Index of colonization success (ICS): **5**
Index of colonization potential (ICP): **9**
Optimum successional age [years]: **5**

Distribution and frequency

Floristic zone: **boreal, northern temperate, southern temperate, submeridional, meridional, subtropical**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Floristic region: **circumpolar**
Distribution range extension along the continentality gradient: **8**
Occurrence frequency in the basic grid mapping cells and quadrants of the basic grid mapping cells: 153
taxon.data.freq_in_quad: 196

Commonness in vegetation plots from the Czech Republic

Occurrence frequency in vegetation plots: **0 %**
Occurrence frequency in vegetation plots with a cover above 5%: **0 %**
Occurrence frequency in vegetation plots with a cover above 25%: **0 %**
Occurrence frequency in vegetation plots with a cover above 50%: **0 %**
Mean percentage cover in vegetation plots: **2 %**
Maximum percentage cover in vegetation plots: **3 %**

Number of habitats with taxon occurrence in the Czech Republic

Number of narrow habitats in which the taxon occurs: **6**
Number of narrow habitats in which the taxon has its optimum: **3**
Number of broad habitats in which the taxon occurs: **3**
Number of broad habitats in which the taxon has its optimum: **3**

Threats and protection

Red List 2017 (national categories): **taxon is not on the Red List**
Red List 2017 (IUCN categories): **LC(NA) – least concern (taxon is not on the Red List)**
Legal protection: **not protected by law**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz