
Kerria japonica

Distribution





Map info

revised records

unrevised records

On the map are not visualized records without the coordinates and records marked as incorrect or doubtful.







Habitus and growth type

Height [m]: **1–2.5**
Growth form: **shrub**
Life form: **nanophanerophyte**
Life strategy (Pierce method based on leaf traits): **CSR**
Life strategy (Pierce method, C-score): **27.9 %**
Life strategy (Pierce method, S-score): **39.1 %**
Life strategy (Pierce method, R-score): **33.1 %**

Leaf

Leaf presence and metamorphosis: **leaves present, not modified**
Leaf arrangement (phyllotaxis): **alternate**
Leaf shape: **simple – entire**
Stipules: **present**
Petiole: **present**
Leaf deciduousness in woody plants: **winter deciduous**
Functional leaf type in woody plants: **broad deciduous or semi-deciduous**

Flower

Flowering period [month]: **May–June**
Flower colour: **yellow**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz



Flower symmetry: **actinomorphic**
Perianth type: **calyx and corolla**
Perianth fusion: **free**
Calyx fusion: **aposepalous**
Inflorescence type: **flores solitarii**

Fruit, seed and dispersal

Fruit type: **dry fruit – head of achenes**
Fruit colour: **brown, black**

Belowground organs and clonality



Position of root buds: **lateral roots**
Role of root buds in life-history of a plant: **regenerative**

Bud bank

Number of buds per shoot at the soil surface (root buds excluded): **0**
Number of buds per shoot at a depth of 0–10 cm (root buds excluded): **5**
Number of buds per shoot at a depth greater than 10 cm (root buds excluded): **5**
Size of the belowground bud bank (root buds excluded): **10**
Depth of the belowground bud bank (root buds exluded) [cm]: **10**
Number of buds per shoot at the soil surface (root buds included): **0**
Number of buds per shoot at a depth of 0–10 cm (root buds included): **20**
Number of buds per shoot at a depth greater than 10 cm (root buds included): **20**
Size of the belowground bud bank (root buds included): **40**
Depth of the belowground bud bank (root buds included) [cm]: **10**

Trophic mode



Parasitism and mycoheterotrophy: **autotrophic**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in the Czech Republic: **cultivated only**

Habitat and sociology

Affinity to the forest environment

Affinity to the forest environment in Thermophyticum: **0 – taxon that does not spontaneously occur in Czech forests**
Affinity to the forest environment in Mesophyticum and Oreophyticum: **0 – taxon that does not spontaneously occur in Czech forests**

Threats and protection

Red List 2017 (national categories): **taxon is not on the Red List**
Red List 2017 (IUCN categories): **LC(NA) – least concern (taxon is not on the Red List)**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Legal protection: **not protected by law**

