
Lathyrus sativus

Distribution





Map info

revised records

unrevised records

On the map are not visualized records without the coordinates and records marked as incorrect or doubtful.



Habitus and growth type

Height [m]: **0.3–0.7**
Growth form: **annual herb**
Life form: **therophyte**
Life strategy: **CR – competitor/ruderal**

Leaf

Leaf presence and metamorphosis: **leaves present, not modified**
Leaf arrangement (phyllotaxis): **alternate**
Leaf shape: **compound – paripinnate**
Stipules: **present**
Petiole: **present**
Leaf life span: **summer green**
Leaf anatomy: **scleromorphic, mesomorphic**

Flower

Flowering period [month]: **May–August**
Flower colour: **white, blue**
Flower symmetry: **zygomorphic**
Perianth type: **calyx and corolla**
Perianth fusion: **free**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Calyx fusion: **synsepalous**
Inflorescence type: **racemus**
Dicliny: **synoecious**
Generative reproduction type: **mixed mating**
Pollination syndrome: **insect-pollination, selfing**

Fruit, seed and dispersal

Fruit type: **dry fruit – legume**
Fruit colour: **brown**
Reproduction type: **only by seed/spores**
Dispersal unit (diaspore): **seed**
Dispersal strategy: **Allium (mainly autochory)**
Myrmecochory: **non-myrmecochorous (b)**

Trophic mode

Parasitism and mycoheterotrophy: **autotrophic**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **symbiosis with rhizobia**

Taxon origin

Origin in the Czech Republic: **archaeophyte**
Invasion status: **casual**
Geographic origin: **anecophyte**
Period of introduction: **Bronze Age (2300–750 BCE)**
Introduction pathway: **intentional – crops, unintentional – anthropogenic**

Habitat and sociology

Affinity to the forest environment

Affinity to the forest environment in Thermophyticum: **0 – taxon that does not spontaneously occur in Czech forests**
Affinity to the forest environment in Mesophyticum and Oreophyticum: **0 – taxon that does not spontaneously occur in Czech forests**

Ecological specialization indices

Ecological specialization index for all vegetation types: **6.4**
Ecological specialization index for non-forest vegetation: **6.4**

Distribution and frequency

Floristic zone: **submeridional, meridional**
Floristic region: **Near East**
Occurrence frequency in the basic grid mapping cells and quadrants of the basic grid mapping cells: 35
taxon.data.freq_in_quad: 40

Commonness in vegetation plots from the Czech Republic

Occurrence frequency in vegetation plots: **0 %**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Occurrence frequency in vegetation plots with a cover above 5%: **25 %**
Occurrence frequency in vegetation plots with a cover above 25%: **0 %**
Occurrence frequency in vegetation plots with a cover above 50%: **0 %**
Mean percentage cover in vegetation plots: **6 %**
Maximum percentage cover in vegetation plots: **18 %**

Threats and protection

Red List 2017 (national categories): **taxon is not on the Red List**
Red List 2017 (IUCN categories): **LC(NA) – least concern (taxon is not on the Red List)**
Legal protection: **not protected by law**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz