
Leucanthemum margaritae

Distribution





Map info

revised records

unrevised records

On the map are not visualized records without the coordinates and records marked as incorrect or doubtful.

Habitus and growth type

Height [m]: **0.25–0.9**
Growth form: **clonal herb**
Life form: **hemicryptophyte**
Life strategy: **CSR – competitor/stress-tolerator/ruderal**

Leaf

Leaf presence and metamorphosis: **leaves present, not modified**
Leaf arrangement (phyllotaxis): **alternate**
Leaf shape: **simple – entire**
Stipules: **absent**
Petiole: **both present and absent**
Leaf life span: **evergreen**
Leaf anatomy: **mesomorphic**

Flower

Flowering period [month]: **June–July**
Flower colour: **white, yellow**
Flower symmetry: **actinomorphic, zygomorphic**
Perianth type: **calyx sometimes absent, corolla present**
Perianth fusion: **fused**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Shape of the sympetalous corolla or syntepalous perianth: **ligulate, tubular**
Calyx fusion: **synsepalous**
Inflorescence type: **anthodium solitarium**
Dicliny: **gynomonoecious**
Generative reproduction type: **mixed mating**
Pollination syndrome: **insect-pollination, selfing**

Fruit, seed and dispersal

Fruit type: **dry fruit – achene/cypsela/samara**
Fruit colour: **brown, black**
Reproduction type: **by seed/spores and vegetatively**
Dispersal unit (diaspore): **fruit, infrutescence or its part**
Dispersal strategy: **Allium (mainly autochory)**
Myrmecochory: **myrmecochorous nv**

Belowground organs and clonality

Shoot metamorphosis: **stolon-like rhizome**
Storage organ: **stolon-like rhizome**

Trophic mode

Parasitism and mycoheterotrophy: **autotrophic**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Karyology

Chromosome number (2n): **54 (52, 53, 55)**
Ploidy level (x): **6**
2C genome size [Mbp]: **22454.13**
1Cx monoploid genome size [Mbp]: **3742.36**
Genomic GC content: **38.6 %**

Taxon origin

Origin in the Czech Republic: **native**

Ecological indicator values

Ellenberg-type indicator values

Light indicator value: **7 – half-light plant, mostly occurring at full light, but also in the shade up to about 30% of diffuse radiation incident in an open area**
Temperature indicator value: **5 – moderate heat indicator, occurring from lowland to montane belt, mainly in submontane-temperate areas**
Moisture indicator value: **5 – indicator of fresh soils, focus on soils of average moisture, missing on wet and on soils that frequently dry out**
Reaction indicator value: **7 – indicator of slightly acidic to slightly basic conditions, never occurring in very acidic conditions**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Nutrient indicator value: **5 – occurring at moderately nutrient-rich sites, and less frequently at poor and rich sites**
Salinity indicator value: **0 – not salt tolerant, glycophyte**

Habitat and sociology

Occurrence in habitats

1 Vegetation of cliffs, screes and walls

1B Siliceous cliffs and block fields: **2 – optimum**

6 Meadows and mesic pastures

6C Pastures and park grasslands: **2 – optimum**

8 Dry grasslands

8A Hercynian dry grasslands on rock outcrops: **2 – optimum**
8B Submediterranean dry grasslands on rock outcrops: **1 – rare occurrence**
8D Broad-leaved dry grasslands: **2 – optimum**

11 Heathlands and scrub

11L Tall mesic and xeric shrub: **1 – rare occurrence**
11N Low xeric scrub: **1 – rare occurrence**

12 Forests

12J Acidophilous thermophilous oak forests: **1 – rare occurrence**
12O Peri-Alpidic pine forests: **1 – rare occurrence**

Affinity to the forest environment

Affinity to the forest environment in Thermophyticum: **0 – taxon that does not spontaneously occur in Czech forests**
Affinity to the forest environment in Mesophyticum and Oreophyticum: **0 – taxon that does not spontaneously occur in Czech forests**

Colonization ability

Index of colonization success (ICS): **1**

Distribution and frequency

Floristic zone: **northern temperate, southern temperate, submeridional**
Floristic region: **Europe**
Elevational belt in the Czech Republic: **colline belt, submontane belt**
Occurrence frequency in the basic grid mapping cells and quadrants of the basic grid mapping cells: 24
taxon.data.freq_in_quad: 38

Number of habitats with taxon occurrence in the Czech Republic

Number of narrow habitats in which the taxon occurs: **9**
Number of narrow habitats in which the taxon has its optimum: **4**
Number of broad habitats in which the taxon occurs: **5**
Number of broad habitats in which the taxon has its optimum: **3**

Threats and protection

Red List 2017 (national categories): **C2r – endangered taxon, rare**
Red List 2017 (IUCN categories): **VU – vulnerable**
Legal protection: **not protected by law**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz