
Ligusticum mutellina

Distribution





Map info

revised records

unrevised records

On the map are not visualized records without the coordinates and records marked as incorrect or doubtful.









Habitus and growth type

Height [m]: **0.1–0.5**
Growth form: **polycarpic perennial non-clonal herb**
Life form: **hemicryptophyte**
Life strategy: **C – competitor**
Life strategy (Pierce method based on leaf traits): **CR**
Life strategy (Pierce method, C-score): **55.3 %**
Life strategy (Pierce method, S-score): **0 %**
Life strategy (Pierce method, R-score): **44.7 %**

Leaf

Leaf presence and metamorphosis: **leaves present, not modified**
Leaf arrangement (phyllotaxis): **alternate**
Leaf shape: **compound – bipinnate, compound – tripinnate**
Stipules: **absent**
Petiole: **both present and absent**
Leaf life span: **evergreen**
Leaf anatomy: **mesomorphic**

Flower

Flowering period [month]: **June–August**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz



Flower colour: **white, pink**
Flower symmetry: **actinomorphic**
Perianth type: **calyx absent, corolla present**
Perianth fusion: **free**
Inflorescence type: **umbella composita**
Dicliny: **andromonoecious**
Generative reproduction type: **facultative allogamy**
Pollination syndrome: **insect-pollination**
Pollinator spectrum: **flies s. l. (hoverflies, meat flies s. l.)**



Fruit, seed and dispersal

Fruit type: **dry fruit – cremocarp**
Fruit colour: **brown**
Reproduction type: **by seed/spores and vegetatively**
Dispersal unit (diaspore): **fruit, infrutescence or its part**
Dispersal strategy: **Allium (mainly autochory)**
Myrmecochory: **non-myrmecochorous (b)**

Belowground organs and clonality

Shoot metamorphosis: **pleiocorm**
Root metamorphosis: **primary storage root, root shoot**
Storage organ: **pleiocorm, primary storage root**
Shoot life span (cyclicity): **dicyclic or polycyclic shoots prevailing**
Branching type of stem-derived organs of clonal growth: **sympodial**
Primary root: **present**
Position of root buds: **primary root**
Role of root buds in life-history of a plant: **additive**

Bud bank

Number of buds per shoot at the soil surface (root buds excluded): **5**
Number of buds per shoot at a depth of 0–10 cm (root buds excluded): **15**
Number of buds per shoot at a depth greater than 10 cm (root buds excluded): **0**
Size of the belowground bud bank (root buds excluded): **20**
Depth of the belowground bud bank (root buds exluded) [cm]: **4**
Number of buds per shoot at the soil surface (root buds included): **5**
Number of buds per shoot at a depth of 0–10 cm (root buds included): **19**
Number of buds per shoot at a depth greater than 10 cm (root buds included): **15**
Size of the belowground bud bank (root buds included): **39**
Depth of the belowground bud bank (root buds included) [cm]: **8**

Trophic mode

Parasitism and mycoheterotrophy: **autotrophic**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Karyology

Chromosome number (2n): **22**
Ploidy level (x): **2**
2C genome size [Mbp]: **1411.44**
1Cx monoploid genome size [Mbp]: **705.72**
Genomic GC content: **37.6 %**

Taxon origin

Origin in the Czech Republic: **native**

Ecological indicator values

Ellenberg-type indicator values

Light indicator value: **7 – half-light plant, mostly occurring at full light, but also in the shade up to about 30% of diffuse radiation incident in an open area**
Temperature indicator value: **3 – cool indicator, occurring mainly in subalpine areas**
Moisture indicator value: **6 – transition between values 5 and 7**
Reaction indicator value: **5 – indicator of moderate acidity, occurring rarely in strongly acidic as well as in neutral to alkaline conditions**
Nutrient indicator value: **3 – occurring at nutrient-poor sites more frequently than at average sites and exceptionally at rich sites**
Salinity indicator value: **0 – not salt tolerant, glycophyte**

Indicator values for disturbance

Whole-community disturbance frequency indicator value: **-0.97**
Herb layer disturbance frequency indicator value: **-0.66**
Whole-community disturbance severity indicator value: **0.18**
Herb layer disturbance severity indicator value: **0.21**
Whole-community structure based disturbance indicator value: **0.47**
Herb layer structure-based disturbance indicator value: **0.57**

Habitat and sociology

Occurrence in habitats

2 Alpine and subalpine grasslands

2A Alpine grasslands on siliceous bedrock: **2 – optimum**
2B Subalpine tall-forb and tall-grass vegetation: **2 – optimum**

5 Vegetation of springs and mires

5C Alpine and subalpine soft-water springs: **1 – rare occurrence**

6 Meadows and mesic pastures

6B Montane mesic meadows: **1 – rare occurrence**

7 Acidophilous grasslands

7A Subalpine and montane acidophilous grasslands: **2 – optimum**
7B Submontane Nardus grasslands: **1 – rare occurrence**

11 Heathlands and scrub

11A Dry lowland to subalpine heathlands: **1 – rare occurrence**
11H Subalpine deciduous scrub: **1 – rare occurrence**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

12 Forests

12R Acidophilous spruce forests: **1 – rare occurrence**

Affinity to the forest environment

Affinity to the forest environment in Thermophyticum: **0 – taxon that does not spontaneously occur in Czech forests**

Affinity to the forest environment in Mesophyticum and Oreophyticum: **0 – taxon that does not spontaneously occur in Czech forests**

Diagnostic taxon

Diagnostic taxon of classes: **AD *Mulgedio-Aconitetea***

Diagnostic taxon of alliances: **ADA *Calamagrostion villosae*, ADD *Adenostylion alliariae***

Diagnostic taxon of associations: **ADA03 *Violo sudeticae-Deschampsietum cespitosae*, ADD01 *Ranunculo platanifolii-Adenostyletum alliariae*, ADD02 *Salicetum lapponum*, TEA02 *Thesio alpini-Nardetum strictae*, TEF03 *Festuco supinae-Vaccinietum myrtilli***

Constant taxon

Constant taxon of associations: **ADA03 *Violo sudeticae-Deschampsietum cespitosae***

Ecological specialization indices

Ecological specialization index for all vegetation types: **5.2**

Ecological specialization index for non-forest vegetation: **5.4**

Colonization ability

Index of colonization success (ICS): **1**

Index of colonization potential (ICP): **1**

Distribution and frequency

Floristic zone: **northern temperate, southern temperate, submeridional**

Floristic region: **Europe**

Continentality degree: **5**

Distribution range extension along the continentality gradient: **3**

Elevational belt in the Czech Republic: **montane belt, subalpine belt**

Occurrence frequency in the basic grid mapping cells and quadrants of the basic grid mapping cells: 37

taxon.data.freq_in_quad: 64

Commonness in vegetation plots from the Czech Republic

Occurrence frequency in vegetation plots: **0.1 %**

Occurrence frequency in vegetation plots with a cover above 5%: **5 %**

Occurrence frequency in vegetation plots with a cover above 25%: **0 %**

Occurrence frequency in vegetation plots with a cover above 50%: **0 %**

Mean percentage cover in vegetation plots: **2.9 %**

Maximum percentage cover in vegetation plots: **13 %**

Number of habitats with taxon occurrence in the Czech Republic

Number of narrow habitats in which the taxon occurs: **9**

Number of narrow habitats in which the taxon has its optimum: **3**

Number of broad habitats in which the taxon occurs: **6**

Number of broad habitats in which the taxon has its optimum: **2**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Threats and protection

Red List 2017 (national categories): **C3 – vulnerable taxon**
Red List 2017 (IUCN categories): **NT – near threatened**
Legal protection: **vulnerable taxon**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz