
Malus domestica



Distribution



Map info

- revised records
- unrevised records

On the map are not visualized records without the coordinates and records marked as incorrect or doubtful.

Habitus and growth type

Height [m]: **6–14**
Growth form: **tree (shrub)**
Life form: **macrophanerophyte (nanophanerophyte)**
Life strategy: **C – competitor**



Leaf

Leaf presence and metamorphosis: **leaves present, not modified**
Leaf arrangement (phyllotaxis): **alternate**
Leaf shape: **simple – entire**
Stipules: **present**
Petiole: **present**
Leaf life span: **summer green**
Leaf deciduousness in woody plants: **winter deciduous**
Leaf anatomy: **mesomorphic**
Functional leaf type in woody plants: **broad deciduous or semi-deciduous**




Flower

Flowering period [month]: **May**
Flower colour: **white**
Flower symmetry: **actinomorphic**



© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz



Perianth type: **calyx and corolla**
Perianth fusion: **free**
Calyx fusion: **hypanthium**
Inflorescence type: **corymbus**
Dicliny: **synoecious**
Generative reproduction type: **allogamy self-incompatibility**
Pollination syndrome: **insect-pollination**

Fruit, seed and dispersal



Fruit type: **fleshy fruit – pome**
Fruit colour: **green, yellow, red**
Reproduction type: **mostly by seed/spores, rarely vegetatively**
Dispersal unit (diaspore): **seed, fruit, infrutescence or its part**
Dispersal strategy: **Cornus (mainly autochory and endozoochory)**
Myrmecochory: **non-myrmecochorous (b)**

Trophic mode



Parasitism and mycoheterotrophy: **autotrophic**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Karyology



Chromosome number (2n): **34**
Ploidy level (x): **2**
2C genome size [Mbp]: **1310.16**
1Cx monoploid genome size [Mbp]: **655.08**
Genomic GC content: **40.4 %**

Taxon origin



Origin in the Czech Republic: **archaeophyte**
Invasion status: **naturalized**
Geographic origin: **anecophyte**
Period of introduction: **Early Middle Ages (550–1200)**
Introduction pathway: **intentional – crops**

Ecological indicator values



Ellenberg-type indicator values

Light indicator value: **6 – transition between values 5 and 7; rarely at less than 20% of diffuse radiation incident in an open area**
Temperature indicator value: **6 – transition between values 5 and 7**
Moisture indicator value: **5 – indicator of fresh soils, focus on soils of average moisture, missing on wet and on soils that frequently dry out**
Reaction indicator value: **7 – indicator of slightly acidic to slightly basic conditions, never occurring in very acidic conditions**
Nutrient indicator value: **5 – occurring at moderately nutrient-rich sites, and less**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

frequently at poor and rich sites
Salinity indicator value: **0 – not salt tolerant, glycophyte**



Habitat and sociology

Occurrence in habitats

1 Vegetation of cliffs, screes and walls

1A Calcareous cliffs: **1 – rare occurrence**
1B Siliceous cliffs and block fields: **1 – rare occurrence**

11 Heathlands and scrub

11L Tall mesic and xeric shrub: **2 – optimum**
11N Low xeric scrub: **1 – rare occurrence**
11R Scrub and pioneer woodland of forests clearings: **2 – optimum**



12 Forests

12B Alluvial forests: **1 – rare occurrence**
12C Oak-hornbeam forests: **1 – rare occurrence**
12D Ravine forests: **1 – rare occurrence**
12H Peri-Alpidic basiphilous thermophilous oak forests: **1 – rare occurrence**
12I Sub-continental thermophilous oak forests: **1 – rare occurrence**
12J Acidophilous thermophilous oak forests: **1 – rare occurrence**
12T Robinia pseudacacia plantations: **1 – rare occurrence**
12U Plantations of broad-leaved non-native trees: **1 – rare occurrence**

13 Anthropogenic vegetation

13E Perennial nitrophilous herbaceous vegetation of mesic sites: **1 – rare occurrence**

Affinity to the forest environment

Affinity to the forest environment in Thermophyticum: **2.2 – taxon occurring partly in the forest, but mainly in open vegetation**
Affinity to the forest environment in Mesophyticum and Oreophyticum: **2.2 – taxon occurring partly in the forest, but mainly in open vegetation**



Colonization ability

Index of colonization success (ICS): **1**

Distribution and frequency

Floristic zone: **submeridional**
Floristic region: **Europe**
Continentality degree: **7**
Distribution range extension along the continentality gradient: **4**
Occurrence frequency in the basic grid mapping cells and quadrants of the basic grid mapping cells: 612
taxon.data.freq_in_quad: 1764



Number of habitats with taxon occurrence in the Czech Republic

Number of narrow habitats in which the taxon occurs: **14**
Number of narrow habitats in which the taxon has its optimum: **2**
Number of broad habitats in which the taxon occurs: **4**
Number of broad habitats in which the taxon has its optimum: **1**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Threats and protection

Red List 2017 (national categories): **taxon is not on the Red List**
Red List 2017 (IUCN categories): **LC(NA) – least concern (taxon is not on the Red List)**
Legal protection: **not protected by law**



