
Melilotus altissimus

Distribution





Map info

revised records

unrevised records

On the map are not visualized records without the coordinates and records marked as incorrect or doubtful.







Habitus and growth type

Height [m]: **0.5–1.5**
Growth form: **monocarpic perennial non-clonal herb**
Life form: **hemicryptophyte**
Life strategy: **CR – competitor/ruderal**
Life strategy (Pierce method based on leaf traits): **S/CSR**
Life strategy (Pierce method, C-score): **18.5 %**
Life strategy (Pierce method, S-score): **49.2 %**
Life strategy (Pierce method, R-score): **32.3 %**

Leaf

Leaf presence and metamorphosis: **leaves present, not modified**
Leaf arrangement (phyllotaxis): **alternate**
Leaf shape: **compound – ternate**
Stipules: **present**
Petiole: **present**
Leaf life span: **summer green**
Leaf anatomy: **mesomorphic**

Flower

Flowering period [month]: **July–September**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz



Flowering phase: **8 Clematis vitalba-Galium sylvaticum (mid-summer)**
Flower colour: **yellow**
Flower symmetry: **zygomorphic**
Perianth type: **calyx and corolla**
Perianth fusion: **free**
Calyx fusion: **synsepalous**
Inflorescence type: **racemus**
Dicliny: **synoecious**
Generative reproduction type: **mixed mating**
Pollination syndrome: **insect-pollination**

Fruit, seed and dispersal

Fruit type: **dry fruit – legume**
Fruit colour: **brown, grey**
Reproduction type: **only by seed/spores**
Dispersal unit (diaspore): **seed, fruit, infrutescence or its part**
Dispersal strategy: **Allium (mainly autochory)**
Myrmecochory: **non-myrmecochorous (b)**

Belowground organs and clonality

Root metamorphosis: **primary storage root**
Storage organ: **primary storage root**
Shoot life span (cyclicity): **dicyclic or polycyclic shoots prevailing**
Primary root: **present**

Bud bank

Number of buds per shoot at the soil surface (root buds excluded): **5**
Number of buds per shoot at a depth of 0–10 cm (root buds excluded): **5**
Number of buds per shoot at a depth greater than 10 cm (root buds excluded): **0**
Size of the belowground bud bank (root buds excluded): **10**
Depth of the belowground bud bank (root buds exluded) [cm]: **3**
Number of buds per shoot at the soil surface (root buds included): **5**
Number of buds per shoot at a depth of 0–10 cm (root buds included): **5**
Number of buds per shoot at a depth greater than 10 cm (root buds included): **0**
Size of the belowground bud bank (root buds included): **10**
Depth of the belowground bud bank (root buds included) [cm]: **3**

Trophic mode

Parasitism and mycoheterotrophy: **autotrophic**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **symbiosis with rhizobia**

Karyology

Chromosome number (2n): **16**
Ploidy level (x): **2**
2C genome size [Mbp]: **2548.99**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

1Cx monoploid genome size [Mbp]: **1274.5**
Genomic GC content: **39 %**

Taxon origin

Origin in the Czech Republic: **native**

Ecological indicator values

Ellenberg-type indicator values

Light indicator value: **8 – light plant, only exceptionally occurring at less than 40% of diffuse radiation incident in an open area**
Temperature indicator value: **6 – transition between values 5 and 7**
Moisture indicator value: **7 – humidity indicator, focus on well moistened, but not wet soils**
Reaction indicator value: **7 – indicator of slightly acidic to slightly basic conditions, never occurring in very acidic conditions**
Nutrient indicator value: **6 – transition between values 5 and 7**
Salinity indicator value: **2 – oligohaline, often on soils with very low salt content**

Habitat and sociology

Occurrence in habitats

4 Wetland and riverine herbaceous vegetation

4B Halophilous reed and sedge beds: **2 – optimum**
4L Nitrophilous herbaceous fringes of lowland rivers: **1 – rare occurrence**

6 Meadows and mesic pastures

6C Pastures and park grasslands: **1 – rare occurrence**
6D Alluvial meadows of lowland rivers: **1 – rare occurrence**
6F Intermittently wet Molinia meadows: **1 – rare occurrence**
6G Vegetation of wet disturbed soils: **1 – rare occurrence**

10 Saline vegetation

10I Inland saline meadows: **1 – rare occurrence**

11 Heathlands and scrub

11J Willow galleries of loamy and sandy river banks: **1 – rare occurrence**

13 Anthropogenic vegetation

13A Annual vegetation of ruderal habitats: **1 – rare occurrence**
13D Perennial thermophilous ruderal vegetation: **1 – rare occurrence**
13E Perennial nitrophilous herbaceous vegetation of mesic sites: **1 – rare occurrence**

Affinity to the forest environment

Affinity to the forest environment in Thermophyticum: **0 – taxon that does not spontaneously occur in Czech forests**
Affinity to the forest environment in Mesophyticum and Oreophyticum: **0 – taxon that does not spontaneously occur in Czech forests**

Colonization ability

Index of colonization success (ICS): **1**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Distribution and frequency

Floristic zone: **northern temperate, southern temperate, submeridional**
Floristic region: **Europe, Western Asia**
Continentality degree: **5**
Distribution range extension along the continentality gradient: **4**
Elevational belt in the Czech Republic: **lowlands, colline belt**
Occurrence frequency in the basic grid mapping cells and quadrants of the basic grid mapping cells: 124
taxon.data.freq_in_quad: 208

Commonness in vegetation plots from the Czech Republic

Occurrence frequency in vegetation plots: **0 %**
Occurrence frequency in vegetation plots with a cover above 5%: **0 %**
Occurrence frequency in vegetation plots with a cover above 25%: **0 %**
Occurrence frequency in vegetation plots with a cover above 50%: **0 %**
Mean percentage cover in vegetation plots: **2 %**
Maximum percentage cover in vegetation plots: **2 %**

Number of habitats with taxon occurrence in the Czech Republic

Number of narrow habitats in which the taxon occurs: **11**
Number of narrow habitats in which the taxon has its optimum: **1**
Number of broad habitats in which the taxon occurs: **5**
Number of broad habitats in which the taxon has its optimum: **1**

Threats and protection

Red List 2017 (national categories): **C3 – vulnerable taxon**
Red List 2017 (IUCN categories): **VU – vulnerable**
Legal protection: **not protected by law**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz