
Montia arvensis

Distribution





Map info

revised records

unrevised records

On the map are not visualized records without the coordinates and records marked as incorrect or doubtful.







Habitus and growth type

Height [m]: **0.02–0.09**
Growth form: **annual herb**
Life form: **therophyte**

Leaf

Leaf presence and metamorphosis: **leaves present, not modified**
Leaf arrangement (phyllotaxis): **opposite**
Leaf shape: **simple – entire**
Stipules: **absent**
Petiole: **absent**

Flower

Flowering period [month]: **April–May**
Flower colour: **white**
Flower symmetry: **zygomorphic**
Perianth type: **homochlamydeous**
Perianth fusion: **fused**
Shape of the sympetalous corolla or syntepalous perianth: **rotate**
Inflorescence type: **cincinnus**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Fruit, seed and dispersal

Fruit type: **dry fruit – capsule**
Fruit colour: **brown**
Dispersal unit (diaspore): **seed**
Dispersal strategy: **Sparganium (mainly autochory and hydrochory)**
Myrmecochory: **myrmecochorous**

Trophic mode

Parasitism and mycoheterotrophy: **autotrophic**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Karyology

Chromosome number (2n): **20**
Ploidy level (x): **2**
2C genome size [Mbp]: **556.3**
1Cx monoploid genome size [Mbp]: **278.15**
Genomic GC content: **37.4 %**

Taxon origin

Origin in the Czech Republic: **native**

Ecological indicator values

Ellenberg-type indicator values

Light indicator value: **8 – light plant, only exceptionally occurring at less than 40% of diffuse radiation incident in an open area**
Temperature indicator value: **4 – transition between values 3 and 5**
Moisture indicator value: **7 – humidity indicator, focus on well moistened, but not wet soils**
Reaction indicator value: **4 – transition between values 3 and 5**
Nutrient indicator value: **3 – occurring at nutrient-poor sites more frequently than at average sites and exceptionally at rich sites**
Salinity indicator value: **0 – not salt tolerant, glycophyte**

Habitat and sociology

Affinity to the forest environment

Affinity to the forest environment in Thermophyticum: **0 – taxon that does not spontaneously occur in Czech forests**
Affinity to the forest environment in Mesophyticum and Oreophyticum: **0 – taxon that does not spontaneously occur in Czech forests**

Distribution and frequency

Elevational belt in the Czech Republic: **colline belt**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Occurrence frequency in the basic grid mapping cells and quadrants of the basic grid mapping cells: 21
taxon.data.freq_in_quad: 26

Threats and protection

Red List 2017 (national categories): **C1t – critically threatened taxon, declining**
Red List 2017 (IUCN categories): **CR – critically endangered**
Legal protection: **not protected by law**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz